

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
15 March 2001 (15.03.2001)

PCT

(10) International Publication Number  
**WO 01/18023 A1**

- (51) International Patent Classification<sup>7</sup>: C07H 21/04, C12N 15/00, C12Q 1/68, A61K 48/00
- (21) International Application Number: PCT/US00/24382
- (22) International Filing Date:  
6 September 2000 (06.09.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
09/392,350 9 September 1999 (09.09.1999) US
- (71) Applicant (for all designated States except US): ISIS PHARMACEUTICALS, INC. [US/US]; 2292 Faraday Avenue, Carlsbad, CA 92008 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): MONIA, Brett, P. [US/US]; 7605 Nueva Castilla Way, La Costa, CA 92009 (US). COWSERT, Lex, M. [US/US]; 3008 Newshire Street, Carlsbad, CA 92008 (US).
- (74) Agents: LICATA, Jane, Massey et al.; Law Offices of Jane Massey Licata, 66 E. Main Street, Marlton, NJ 08053 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:  
— With international search report.
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 01/18023 A1

(54) Title: ANTISENSE MODULATION OF PI3 KINASE P110 BETA EXPRESSION

(57) Abstract: Antisense compounds, compositions and methods are provided for modulating the expression of PI3 kinase p110 beta. The compositions comprise antisense compounds, particularly antisense oligonucleotides, targeted to nucleic acids encoding PI3 kinase p110 beta. Methods of using these compounds for modulation of PI3 kinase p110 beta expression and for treatment of diseases associated with expression of PI3 kinase p110 beta are provided.



-1-

## ANTISENSE MODULATION OF PI3 KINASE P110 BETA EXPRESSION

## FIELD OF THE INVENTION

The present invention provides compositions and  
5 methods for modulating the expression of PI3 kinase p110  
beta. In particular, this invention relates to antisense  
compounds, particularly oligonucleotides, specifically  
hybridizable with nucleic acids encoding human PI3 kinase  
p110 beta. Such oligonucleotides have been shown to  
10 modulate the expression of PI3 kinase p110 beta.

## BACKGROUND OF THE INVENTION

Many growth factors and hormones such as nerve growth  
factor (NGF), platelet derived growth factor (PDGF),  
epidermal growth factor (EGF) and insulin mediate their  
15 signals through interactions with cell surface tyrosine  
kinase receptors. The transduction of extracellular  
signals across the membrane, initiated by ligand binding,  
leads to the propagation of multiple signaling events which  
ultimately control target biochemical pathways within the  
20 cell.

The phosphatidylinositol 3-kinases (PI3 kinases or  
PI3Ks) represent a ubiquitous family of heterodimeric lipid  
kinases that are found in association with the cytoplasmic  
domain of hormone and growth factor receptors and oncogene  
25 products. PI3Ks act as downstream effectors of these  
receptors, are recruited upon receptor stimulation and  
mediate the activation of second messenger signaling  
pathways through the production of phosphorylated  
derivatives of inositol. PI3Ks phosphorylate  
30 phosphatidylinositol (PtdIns) at the 3'-hydroxyl of the  
inositol ring and substrates include PtdIns,  
PtIns(4)phosphate, PtdIns(4,5)bisphosphate and  
PtdIns(3,4,5) triphosphate with the major product being  
PtdIns(3,4,5)triphosphate (Fry, *Biochim. Biophys. Acta.*,  
35 1994, 1226, 237-268).

-2-

PI3Ks have been implicated in many cellular activities including growth factor mediated cell transformation, mitogenesis, protein trafficking, cell survival and proliferation, DNA synthesis, apoptosis, neurite outgrowth and insulin-stimulated glucose transport (Fry, *Biochim. Biophys. Acta.*, 1994, 1226, 237-268).

The PI3 kinase enzyme heterodimers consist of a 110 kD (p110) catalytic subunit associated with an 85 kD (p85) regulatory subunit and it is through the SH2 domains of the p85 subunit that the enzyme associates with the membrane-bound receptors. Multiple isoforms of the p110 catalytic subunit have been reported and they have been divided into three classes, IA and IB, II, and III. Class IA includes the alpha, beta and delta isoforms all of which bind to p85 and associate with receptor tyrosine kinases (Hu et al., *Mol. Cell. Biol.*, 1993, 13, 7677-7688; Stirdivant et al., *Bioorg. Med. Chem.*, 1997, 5, 65-74; Vanhaesebroeck et al., *Proc. Natl. Acad. Sci. U. S. A.*, 1997, 94, 4330-4335).

While the alpha and beta isoforms are ubiquitously expressed, the delta isoform is found only in white blood cells (Vanhaesebroeck et al., *Proc. Natl. Acad. Sci. U. S. A.*, 1997, 94, 4330-4335). Class IB includes the gamma isoform of p110 which exhibits no p85 binding, is regulated by the  $\alpha$  and  $\beta\gamma$  subunits of G-proteins, and contains unique C2 domains (Stoyanov et al., *Science*, 1995, 269, 690-693). Classes II and III contain the cpk and Vps34p isoforms, respectively (Molz et al., *J. Biol. Chem.*, 1996, 271, 13892-13899; Schu et al., *Science*, 1993, 260, 88-91).

PI3 kinase p110 beta was first isolated by screening a human cDNA library with a probe derived from regions conserved between the bovine and yeast proteins. In these characterization studies, the PI3 kinase p110 beta isoform was found to exhibit 42% identity to the bovine protein and

-3-

to be ubiquitously expressed in all mouse tissues tested (Hu et al., *Mol. Cell. Biol.*, 1993, 13, 7677-7688).

A role for PI3 kinase p110 beta in several signaling pathways including those involving G-proteins, MAP kinases, and protein kinase C, has been demonstrated in the literature. Using the human colon carcinoma cell line SW-480, which only expresses the p110 beta isoform, bradykinin stimulation led to increased formation of PI3 kinase lipid products and enhanced translocation of the epsilon isoform of protein kinase C to the cell membrane. These effects were inhibited by wortmannin, a PI3 kinase specific inhibitor. In addition, both bradykinin-induced DNA synthesis and activation of MAP kinase were abolished by wortmannin and LY294002, another PI3 kinase inhibitor (Graness et al., *J. Biol. Chem.*, 1998, 273, 32016-32022).

It has also been demonstrated that nitrosative and oxidative stressors can trigger ras- and PI3 kinase-regulated events in the cell. Studies of redox signaling in human T cells showed that only the beta and delta isoforms of PI3 kinase p110 were recruited to the membrane upon free radical generation from nitric oxide donors (Deora et al., *J. Biol. Chem.*, 1998, 273, 29923-29928). These findings suggest a role for PI3 kinase beta in inflammatory conditions such as rheumatoid arthritis and asthma.

Currently, there are no known therapeutic agents which effectively inhibit the synthesis of PI3 kinase p110 beta.

To date, strategies aimed at inhibiting PI3 kinase p110 beta function have involved the use of antibodies and chemical inhibitors.

Several chemically distinct inhibitors for PI3 kinases are reported in the literature. These include wortmannin, a fungal metabolite (Ui et al., *Trends Biochem. Sci.*, 1995, 20, 303-307); demethoxyviridin, an antifungal

-4-

agent (Woscholski et al., *FEBS Lett.*, 1994, 342, 109-114) and quercetin and LY294002, two related chromones (Vlahos et al., *J. Biol. Chem.*, 1994, 269, 5241-5248). However, these strategies are untested as therapeutic protocols as well as being non-specific to the PI3 kinase p110 beta isoform. Consequently, there remains a long felt need for additional agents capable of effectively inhibiting PI3 kinase p110 beta function.

Antisense technology is emerging as an effective means for reducing the expression of specific gene products and may therefore prove to be uniquely useful in a number of therapeutic, diagnostic, and research applications for the modulation of PI3 kinase p110 beta expression.

#### SUMMARY OF THE INVENTION

The present invention is directed to antisense compounds, particularly oligonucleotides, which are targeted to a nucleic acid encoding PI3 kinase p110 beta, and which modulate the expression of PI3 kinase p110 beta. Pharmaceutical and other compositions comprising the antisense compounds of the invention are also provided. Further provided are methods of modulating the expression of PI3 kinase p110 beta in cells or tissues comprising contacting said cells or tissues with one or more of the antisense compounds or compositions of the invention. Further provided are methods of treating an animal, particularly a human, suspected of having or being prone to a disease or condition associated with expression of PI3 kinase p110 beta by administering a therapeutically or prophylactically effective amount of one or more of the antisense compounds or compositions of the invention.

#### DETAILED DESCRIPTION OF THE INVENTION

The present invention employs oligomeric antisense compounds, particularly oligonucleotides, for use in modulating the function of nucleic acid molecules encoding PI3 kinase p110 beta, ultimately modulating the amount of

-5-

PI3 kinase p110 beta produced. This is accomplished by providing antisense compounds which specifically hybridize with one or more nucleic acids encoding PI3 kinase p110 beta. As used herein, the terms "target nucleic acid" and

5 "nucleic acid encoding PI3 kinase p110 beta" encompass DNA encoding PI3 kinase p110 beta, RNA (including pre-mRNA and mRNA) transcribed from such DNA, and also cDNA derived from such RNA. The specific hybridization of an oligomeric compound with its target nucleic acid interferes with the

10 normal function of the nucleic acid. This modulation of function of a target nucleic acid by compounds which specifically hybridize to it is generally referred to as "antisense". The functions of DNA to be interfered with include replication and transcription. The functions of

15 RNA to be interfered with include all vital functions such as, for example, translocation of the RNA to the site of protein translation, translation of protein from the RNA, splicing of the RNA to yield one or more mRNA species, and catalytic activity which may be engaged in or facilitated

20 by the RNA. The overall effect of such interference with target nucleic acid function is modulation of the expression of PI3 kinase p110 beta. In the context of the present invention, "modulation" means either an increase (stimulation) or a decrease (inhibition) in the expression

25 of a gene. In the context of the present invention, inhibition is the preferred form of modulation of gene expression and mRNA is a preferred target.

It is preferred to target specific nucleic acids for antisense. "Targeting" an antisense compound to a

30 particular nucleic acid, in the context of this invention, is a multistep process. The process usually begins with the identification of a nucleic acid sequence whose function is to be modulated. This may be, for example, a cellular gene (or mRNA transcribed from the gene) whose

35 expression is associated with a particular disorder or

-6-

disease state, or a nucleic acid molecule from an infectious agent. In the present invention, the target is a nucleic acid molecule encoding PI3 kinase p110 beta. The targeting process also includes determination of a site or  
5 sites within this gene for the antisense interaction to occur such that the desired effect, e.g., detection or modulation of expression of the protein, will result. Within the context of the present invention, a preferred intragenic site is the region encompassing the translation  
10 initiation or termination codon of the open reading frame (ORF) of the gene. Since, as is known in the art, the translation initiation codon is typically 5'-AUG (in transcribed mRNA molecules; 5'-ATG in the corresponding DNA molecule), the translation initiation codon is also  
15 referred to as the "AUG codon," the "start codon" or the "AUG start codon". A minority of genes have a translation initiation codon having the RNA sequence 5'-GUG, 5'-UUG or 5'-CUG, and 5'-AUA, 5'-ACG and 5'-CUG have been shown to function in vivo. Thus, the terms "translation initiation  
20 codon" and "start codon" can encompass many codon sequences, even though the initiator amino acid in each instance is typically methionine (in eukaryotes) or formylmethionine (in prokaryotes). It is also known in the  
~~art that eukaryotic and prokaryotic genes may have two or~~  
25 more alternative start codons, any one of which may be preferentially utilized for translation initiation in a particular cell type or tissue, or under a particular set of conditions. In the context of the invention, "start codon" and "translation initiation codon" refer to the  
30 codon or codons that are used in vivo to initiate translation of an mRNA molecule transcribed from a gene encoding PI3 kinase p110 beta, regardless of the sequence(s) of such codons.

It is also known in the art that a translation  
35 termination codon (or "stop codon") of a gene may have one



-7-

of three sequences, i.e., 5'-UAA, 5'-UAG and 5'-UGA (the corresponding DNA sequences are 5'-TAA, 5'-TAG and 5'-TGA, respectively). The terms "start codon region" and "translation initiation codon region" refer to a portion of  
5 such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation initiation codon. Similarly, the terms "stop codon region" and "translation termination codon region" refer to a portion of such an mRNA or gene  
10 that encompasses from about 25 to about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation termination codon.

The open reading frame (ORF) or "coding region," which is known in the art to refer to the region between  
15 the translation initiation codon and the translation termination codon, is also a region which may be targeted effectively. Other target regions include the 5' untranslated region (5'UTR), known in the art to refer to the portion of an mRNA in the 5' direction from the  
20 translation initiation codon, and thus including nucleotides between the 5' cap site and the translation initiation codon of an mRNA or corresponding nucleotides on the gene, and the 3' untranslated region (3'UTR), known in the art to refer to the portion of an mRNA in the 3'  
25 direction from the translation termination codon, and thus including nucleotides between the translation termination codon and 3' end of an mRNA or corresponding nucleotides on the gene. The 5' cap of an mRNA comprises an N7-methylated guanosine residue joined to the 5'-most residue of the mRNA  
30 via a 5'-5' triphosphate linkage. The 5' cap region of an mRNA is considered to include the 5' cap structure itself as well as the first 50 nucleotides adjacent to the cap. The 5' cap region may also be a preferred target region.

Although some eukaryotic mRNA transcripts are  
35 directly translated, many contain one or more regions,

-8-

known as "introns," which are excised from a transcript before it is translated. The remaining (and therefore translated) regions are known as "exons" and are spliced together to form a continuous mRNA sequence. mRNA splice sites, i.e., intron-exon junctions, may also be preferred target regions, and are particularly useful in situations where aberrant splicing is implicated in disease, or where an overproduction of a particular mRNA splice product is implicated in disease. Aberrant fusion junctions due to rearrangements or deletions are also preferred targets. It has also been found that introns can also be effective, and therefore preferred, target regions for antisense compounds targeted, for example, to DNA or pre-mRNA.

Once one or more target sites have been identified, oligonucleotides are chosen which are sufficiently complementary to the target, i.e., hybridize sufficiently well and with sufficient specificity, to give the desired effect.

In the context of this invention, "hybridization" means hydrogen bonding, which may be Watson-Crick, Hoogsteen or reversed Hoogsteen hydrogen bonding, between complementary nucleoside or nucleotide bases. For example, adenine and thymine are complementary nucleobases which

~~pair through the formation of hydrogen bonds.~~

"Complementary," as used herein, refers to the capacity for precise pairing between two nucleotides. For example, if a nucleotide at a certain position of an oligonucleotide is capable of hydrogen bonding with a nucleotide at the same position of a DNA or RNA molecule, then the oligonucleotide and the DNA or RNA are considered to be complementary to each other at that position. The oligonucleotide and the DNA or RNA are complementary to each other when a sufficient number of corresponding positions in each molecule are occupied by nucleotides which can hydrogen bond with each other. Thus, "specifically hybridizable"

-9-

and "complementary" are terms which are used to indicate a sufficient degree of complementarity or precise pairing such that stable and specific binding occurs between the oligonucleotide and the DNA or RNA target. It is understood in the art that the sequence of an antisense compound need not be 100% complementary to that of its target nucleic acid to be specifically hybridizable. An antisense compound is specifically hybridizable when binding of the compound to the target DNA or RNA molecule interferes with the normal function of the target DNA or RNA to cause a loss of utility, and there is a sufficient degree of complementarity to avoid non-specific binding of the antisense compound to non-target sequences under conditions in which specific binding is desired, i.e., under physiological conditions in the case of in vivo assays or therapeutic treatment, and in the case of in vitro assays, under conditions in which the assays are performed.

Antisense compounds are commonly used as research reagents and diagnostics. For example, antisense oligonucleotides, which are able to inhibit gene expression with exquisite specificity, are often used by those of ordinary skill to elucidate the function of particular genes. Antisense compounds are also used, for example, to distinguish between functions of various members of a biological pathway. Antisense modulation has, therefore, been harnessed for research use.

The specificity and sensitivity of antisense is also harnessed by those of skill in the art for therapeutic uses. Antisense oligonucleotides have been employed as therapeutic moieties in the treatment of disease states in animals and man. Antisense oligonucleotides have been safely and effectively administered to humans and numerous clinical trials are presently underway. It is thus established that oligonucleotides can be useful therapeutic

-10-

modalities that can be configured to be useful in treatment regimes for treatment of cells, tissues and animals, especially humans.

In the context of this invention, the term

5 "oligonucleotide" refers to an oligomer or polymer of ribonucleic acid (RNA) or deoxyribonucleic acid (DNA) or mimetics thereof. This term includes oligonucleotides composed of naturally-occurring nucleobases, sugars and covalent internucleoside (backbone) linkages as well as  
10 oligonucleotides having non-naturally-occurring portions which function similarly. Such modified or substituted oligonucleotides are often preferred over native forms because of desirable properties such as, for example, enhanced cellular uptake, enhanced affinity for nucleic  
15 acid target and increased stability in the presence of nucleases.

While antisense oligonucleotides are a preferred form of antisense compound, the present invention comprehends other oligomeric antisense compounds, including but not  
20 limited to oligonucleotide mimetics such as are described below. The antisense compounds in accordance with this invention preferably comprise from about 8 to about 30 nucleobases (i.e. from about 8 to about 30 linked  

---

nucleosides). Particularly preferred antisense compounds  
25 are antisense oligonucleotides, even more preferably those comprising from about 12 to about 25 nucleobases. As is known in the art, a nucleoside is a base-sugar combination. The base portion of the nucleoside is normally a heterocyclic base. The two most common classes of such  
30 heterocyclic bases are the purines and the pyrimidines. Nucleotides are nucleosides that further include a phosphate group covalently linked to the sugar portion of the nucleoside. For those nucleosides that include a pentofuranosyl sugar, the phosphate group can be linked to  
35 either the 2', 3' or 5' hydroxyl moiety of the sugar. In

-11-

forming oligonucleotides, the phosphate groups covalently link adjacent nucleosides to one another to form a linear polymeric compound. In turn the respective ends of this linear polymeric structure can be further joined to form a circular structure, however, open linear structures are generally preferred. Within the oligonucleotide structure, the phosphate groups are commonly referred to as forming the internucleoside backbone of the oligonucleotide. The normal linkage or backbone of RNA and DNA is a 3' to 5' phosphodiester linkage.

Specific examples of preferred antisense compounds useful in this invention include oligonucleotides containing modified backbones or non-natural internucleoside linkages. As defined in this specification, oligonucleotides having modified backbones include those that retain a phosphorus atom in the backbone and those that do not have a phosphorus atom in the backbone. For the purposes of this specification, and as sometimes referenced in the art, modified oligonucleotides that do not have a phosphorus atom in their internucleoside backbone can also be considered to be oligonucleosides.

Preferred modified oligonucleotide backbones include, for example, phosphorothioates, chiral phosphorothioates, phosphorodithioates, phosphotriesters, aminoalkyl-phosphotriesters, methyl and other alkyl phosphonates including 3'-alkylene phosphonates and chiral phosphonates, phosphinates, phosphoramidates including 3'-amino phosphoramidate and aminoalkylphosphoramidates, thionophosphoramidates, thionoalkylphosphonates, thionoalkylphosphotriesters, and boranophosphates having normal 3'-5' linkages, 2'-5' linked analogs of these, and those having inverted polarity wherein the adjacent pairs of nucleoside units are linked 3'-5' to 5'-3' or 2'-5' to 5'-2'. Various salts, mixed salts and free acid forms are also included.

-12-

Representative United States patents that teach the preparation of the above phosphorus-containing linkages include, but are not limited to, U.S.: 3,687,808; 4,469,863; 4,476,301; 5,023,243; 5,177,196; 5,188,897; 5,264,423; 5,276,019; 5,278,302; 5,286,717; 5,321,131; 5,399,676; 5,405,939; 5,453,496; 5,455,233; 5,466,677; 5,476,925; 5,519,126; 5,536,821; 5,541,306; 5,550,111; 5,563,253; 5,571,799; 5,587,361; and 5,625,050, certain of which are commonly owned with this application, and each of which is herein incorporated by reference.

Preferred modified oligonucleotide backbones that do not include a phosphorus atom therein have backbones that are formed by short chain alkyl or cycloalkyl internucleoside linkages, mixed heteroatom and alkyl or cycloalkyl internucleoside linkages, or one or more short chain heteroatomic or heterocyclic internucleoside linkages. These include those having morpholino linkages (formed in part from the sugar portion of a nucleoside); siloxane backbones; sulfide, sulfoxide and sulfone backbones; formacetyl and thioformacetyl backbones; methylene formacetyl and thioformacetyl backbones; alkene containing backbones; sulfamate backbones; methyleneimino and methylenehydrazino backbones; sulfonate and sulfonamide backbones, amide backbones, and others having mixed N, O, S and CH<sub>2</sub> component parts.

Representative United States patents that teach the preparation of the above oligonucleosides include, but are not limited to, U.S.: 5,034,506; 5,166,315; 5,185,444; 5,214,134; 5,216,141; 5,235,033; 5,264,562; 5,264,564; 5,405,938; 5,434,257; 5,466,677; 5,470,967; 5,489,677; 5,541,307; 5,561,225; 5,596,086; 5,602,240; 5,610,289; 5,602,240; 5,608,046; 5,610,289; 5,618,704; 5,623,070; 5,663,312; 5,633,360; 5,677,437; and 5,677,439, certain of which are commonly owned with this application, and each of which is herein incorporated by reference.

-13-

In other preferred oligonucleotide mimetics, both the sugar and the internucleoside linkage, i.e., the backbone, of the nucleotide units are replaced with novel groups. The base units are maintained for hybridization with an appropriate nucleic acid target compound. One such oligomeric compound, an oligonucleotide mimetic that has been shown to have excellent hybridization properties, is referred to as a peptide nucleic acid (PNA). In PNA compounds, the sugar-backbone of an oligonucleotide is replaced with an amide containing backbone, in particular an aminoethylglycine backbone. The nucleobases are retained and are bound directly or indirectly to aza nitrogen atoms of the amide portion of the backbone. Representative United States patents that teach the preparation of PNA compounds include, but are not limited to, U.S.: 5,539,082; 5,714,331; and 5,719,262, each of which is herein incorporated by reference. Further teaching of PNA compounds can be found in Nielsen et al., *Science*, 1991, 254, 1497-1500.

Most preferred embodiments of the invention are oligonucleotides with phosphorothioate backbones and oligonucleosides with heteroatom backbones, and in particular  $-\text{CH}_2-\text{NH}-\text{O}-\text{CH}_2-$ ,  $-\text{CH}_2-\text{N}(\text{CH}_3)-\text{O}-\text{CH}_2-$  [known as a methylene (methylimino) or MMI backbone],  $-\text{CH}_2-\text{O}-\text{N}(\text{CH}_3)-\text{CH}_2-$ ,  $-\text{CH}_2-\text{N}(\text{CH}_3)-\text{N}(\text{CH}_3)-\text{CH}_2-$  and  $-\text{O}-\text{N}(\text{CH}_3)-\text{CH}_2-\text{CH}_2-$  [wherein the native phosphodiester backbone is represented as  $-\text{O}-\text{P}-\text{O}-\text{CH}_2-$ ] of the above referenced U.S. patent 5,489,677, and the amide backbones of the above referenced U.S. patent 5,602,240. Also preferred are oligonucleotides having morpholino backbone structures of the above-referenced U.S. patent 5,034,506.

Modified oligonucleotides may also contain one or more substituted sugar moieties. Preferred oligonucleotides comprise one of the following at the 2' position: OH; F; O-, S-, or N-alkyl; O-, S-, or N-alkenyl;

O-, S- or N-alkynyl; or O-alkyl-O-alkyl, wherein the alkyl, alkenyl and alkynyl may be substituted or unsubstituted C<sub>1</sub> to C<sub>10</sub> alkyl or C<sub>2</sub> to C<sub>10</sub> alkenyl and alkynyl. Particularly preferred are O[(CH<sub>2</sub>)<sub>n</sub>O]<sub>m</sub>CH<sub>3</sub>, O(CH<sub>2</sub>)<sub>n</sub>OCH<sub>3</sub>, O(CH<sub>2</sub>)<sub>n</sub>NH<sub>2</sub>,  
5 O(CH<sub>2</sub>)<sub>n</sub>CH<sub>3</sub>, O(CH<sub>2</sub>)<sub>n</sub>ONH<sub>2</sub>, and O(CH<sub>2</sub>)<sub>n</sub>ON[(CH<sub>2</sub>)<sub>n</sub>CH<sub>3</sub>]<sub>2</sub>, where n and m are from 1 to about 10. Other preferred oligonucleotides comprise one of the following at the 2' position: C<sub>1</sub> to C<sub>10</sub> lower alkyl, substituted lower alkyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH<sub>3</sub>, OCN, Cl, Br, CN, CF<sub>3</sub>, OCF<sub>3</sub>,  
10 SOCH<sub>3</sub>, SO<sub>2</sub>CH<sub>3</sub>, ONO<sub>2</sub>, NO<sub>2</sub>, N<sub>3</sub>, NH<sub>2</sub>, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, an RNA cleaving group, a reporter group, an intercalator, a group for improving the pharmacokinetic properties of an oligonucleotide, or a group for improving  
15 the pharmacodynamic properties of an oligonucleotide, and other substituents having similar properties. A preferred modification includes 2'-methoxyethoxy (2'-O-CH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, also known as 2'-O-(2-methoxyethyl) or 2'-MOE) (Martin et al., *Helv. Chim. Acta*, 1995, 78, 486-504) i.e., an alkoxyalkoxy  
20 group. A further preferred modification includes 2'-dimethylaminoethoxyethoxy, i.e., a O(CH<sub>2</sub>)<sub>2</sub>ON(CH<sub>3</sub>)<sub>2</sub> group, also known as 2'-DMAOE, as described in examples hereinbelow, and 2'-dimethylaminoethoxyethoxy (also known in the art as  
~~2'-O-dimethylaminoethoxyethyl or 2'-DMAEOE~~), i.e., 2'-O-CH<sub>2</sub>-  
25 O-CH<sub>2</sub>-N(CH<sub>3</sub>)<sub>2</sub>, also described in examples hereinbelow.

Other preferred modifications include 2'-methoxy (2'-O-CH<sub>3</sub>), 2'-aminopropoxy (2'-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>) and 2'-fluoro (2'-F). Similar modifications may also be made at other positions on the oligonucleotide, particularly the 3'  
30 position of the sugar on the 3' terminal nucleotide or in 2'-5' linked oligonucleotides and the 5' position of 5' terminal nucleotide. Oligonucleotides may also have sugar mimetics such as cyclobutyl moieties in place of the pentofuranosyl sugar. Representative United States patents  
35 that teach the preparation of such modified sugar



-15-

structures include, but are not limited to, U.S.:

4,981,957; 5,118,800; 5,319,080; 5,359,044; 5,393,878;  
5,446,137; 5,466,786; 5,514,785; 5,519,134; 5,567,811;  
5,576,427; 5,591,722; 5,597,909; 5,610,300; 5,627,053;  
5 5,639,873; 5,646,265; 5,658,873; 5,670,633; and 5,700,920,  
certain of which are commonly owned with the instant  
application, and each of which is herein incorporated by  
reference in its entirety.

Oligonucleotides may also include nucleobase (often  
10 referred to in the art simply as "base") modifications or  
substitutions. As used herein, "unmodified" or "natural"  
nucleobases include the purine bases adenine (A) and  
guanine (G), and the pyrimidine bases thymine (T), cytosine  
(C) and uracil (U). Modified nucleobases include other  
15 synthetic and natural nucleobases such as 5-methylcytosine  
(5-me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine,  
2-aminoadenine, 6-methyl and other alkyl derivatives of  
adenine and guanine, 2-propyl and other alkyl derivatives  
of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-  
20 thiocytosine, 5-halouracil and cytosine, 5-propynyl uracil  
and cytosine, 6-azo uracil, cytosine and thymine, 5-uracil  
(pseudouracil), 4-thiouracil, 8-halo, 8-amino, 8-thiol, 8-  
thioalkyl, 8-hydroxyl and other 8-substituted adenines and  
guanines, 5-halo particularly 5-bromo, 5-trifluoromethyl  
25 and other 5-substituted uracils and cytosines, 7-  
methylguanine and 7-methyladenine, 8-azaguanine and 8-  
azaadenine, 7-deazaguanine and 7-deazaadenine and 3-  
deazaguanine and 3-deazaadenine. Further nucleobases  
include those disclosed in United States Patent No.  
30 3,687,808, those disclosed in *The Concise Encyclopedia Of  
Polymer Science And Engineering*, pages 858-859, Kroschwitz,  
J.I., ed. John Wiley & Sons, 1990, those disclosed by  
Englich et al., *Angewandte Chemie*, International Edition,  
1991, 30, 613, and those disclosed by Sanghvi, Y.S.,  
35 Chapter 15, *Antisense Research and Applications*, pages 289-

-16-

302, Crooke, S.T. and Lebleu, B. , ed., CRC Press, 1993. Certain of these nucleobases are particularly useful for increasing the binding affinity of the oligomeric compounds of the invention. These include 5-substituted pyrimidines, 5 6-azapyrimidines and N-2, N-6 and O-6 substituted purines, including 2-aminopropyladenine, 5-propynyluracil and 5-propynylcytosine. 5-methylcytosine substitutions have been shown to increase nucleic acid duplex stability by 0.6-1.2°C (Sanghvi, Y.S., Crooke, S.T. and Lebleu, B., eds.,  
10 *Antisense Research and Applications*, CRC Press, Boca Raton, 1993, pp. 276-278) and are presently preferred base substitutions, even more particularly when combined with 2'-O-methoxyethyl sugar modifications.

Representative United States patents that teach the  
15 preparation of certain of the above noted modified nucleobases as well as other modified nucleobases include, but are not limited to, the above noted U.S. 3,687,808, as well as U.S.: 4,845,205; 5,130,302; 5,134,066; 5,175,273; 5,367,066; 5,432,272; 5,457,187; 5,459,255; 5,484,908;  
20 5,502,177; 5,525,711; 5,552,540; 5,587,469; 5,594,121, 5,596,091; 5,614,617; and 5,681,941, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference, and United  
States patent 5,750,692, which is commonly owned with the  
25 instant application and also herein incorporated by reference.

Another modification of the oligonucleotides of the invention involves chemically linking to the  
oligonucleotide one or more moieties or conjugates which  
30 enhance the activity, cellular distribution or cellular uptake of the oligonucleotide. Such moieties include but are not limited to lipid moieties such as a cholesterol moiety (Letsinger et al., *Proc. Natl. Acad. Sci. USA*, 1989, 86, 6553-6556), cholic acid (Manoharan et al., *Bioorg. Med.*

-17-

Chem. Lett., 1994, 4, 1053-1060), a thioether, e.g., hexyl-S-tritylthiol (Manoharan et al., Ann. N.Y. Acad. Sci., 1992, 660, 306-309; Manoharan et al., Bioorg. Med. Chem. Lett., 1993, 3, 2765-2770), a thiocholesterol (Oberhauser et al., Nucl. Acids Res., 1992, 20, 533-538), an aliphatic chain, e.g., dodecandiol or undecyl residues (Saison-Behmoaras et al., EMBO J., 1991, 10, 1111-1118; Kabanov et al., FEBS Lett., 1990, 259, 327-330; Svinarchuk et al., Biochimie, 1993, 75, 49-54), a phospholipid, e.g., di-hexadecyl-rac-glycerol or triethylammonium 1,2-di-O-hexadecyl-rac-glycero-3-H-phosphonate (Manoharan et al., Tetrahedron Lett., 1995, 36, 3651-3654; Shea et al., Nucl. Acids Res., 1990, 18, 3777-3783), a polyamine or a polyethylene glycol chain (Manoharan et al., Nucleosides & Nucleotides, 1995, 14, 969-973), or adamantane acetic acid (Manoharan et al., Tetrahedron Lett., 1995, 36, 3651-3654), a palmityl moiety (Mishra et al., Biochim. Biophys. Acta, 1995, 1264, 229-237), or an octadecylamine or hexylamino-carbonyl-oxycholesterol moiety (Crooke et al., J. Pharmacol. Exp. Ther., 1996, 277, 923-937).

Representative United States patents that teach the preparation of such oligonucleotide conjugates include, but are not limited to, U.S.: 4,828,979; 4,948,882; 5,218,105; 5,525,465; 5,541,313; 5,545,730; 5,552,538; 5,578,717; 5,580,731; 5,580,731; 5,591,584; 5,109,124; 5,118,802; 5,138,045; 5,414,077; 5,486,603; 5,512,439; 5,578,718; 5,608,046; 4,587,044; 4,605,735; 4,667,025; 4,762,779; 4,789,737; 4,824,941; 4,835,263; 4,876,335; 4,904,582; 4,958,013; 5,082,830; 5,112,963; 5,214,136; 5,082,830; 5,112,963; 5,214,136; 5,245,022; 5,254,469; 5,258,506; 5,262,536; 5,272,250; 5,292,873; 5,317,098; 5,371,241; 5,391,723; 5,416,203; 5,451,463; 5,510,475; 5,512,667; 5,514,785; 5,565,552; 5,567,810; 5,574,142; 5,585,481; 5,587,371; 5,595,726; 5,597,696; 5,599,923; 5,599,928 and

-18-

5,688,941, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference.

It is not necessary for all positions in a given compound to be uniformly modified, and in fact more than one of the aforementioned modifications may be incorporated in a single compound or even at a single nucleoside within an oligonucleotide. The present invention also includes antisense compounds which are chimeric compounds.

10 "Chimeric" antisense compounds or "chimeras," in the context of this invention, are antisense compounds, particularly oligonucleotides, which contain two or more chemically distinct regions, each made up of at least one monomer unit, i.e., a nucleotide in the case of an

15 oligonucleotide compound. These oligonucleotides typically contain at least one region wherein the oligonucleotide is modified so as to confer upon the oligonucleotide increased resistance to nuclease degradation, increased cellular uptake, and/or increased binding affinity for the target

20 nucleic acid. An additional region of the oligonucleotide may serve as a substrate for enzymes capable of cleaving RNA:DNA or RNA:RNA hybrids. By way of example, RNase H is a cellular endonuclease which cleaves the RNA strand of an

---

25 in cleavage of the RNA target, thereby greatly enhancing the efficiency of oligonucleotide inhibition of gene expression. Consequently, comparable results can often be obtained with shorter oligonucleotides when chimeric oligonucleotides are used, compared to phosphorothioate

30 deoxyoligonucleotides hybridizing to the same target region. Cleavage of the RNA target can be routinely detected by gel electrophoresis and, if necessary, associated nucleic acid hybridization techniques known in the art.

-19-

Chimeric antisense compounds of the invention may be formed as composite structures of two or more oligonucleotides, modified oligonucleotides, oligonucleosides and/or oligonucleotide mimetics as described above. Such compounds have also been referred to in the art as hybrids or gapmers. Representative United States patents that teach the preparation of such hybrid structures include, but are not limited to, U.S.: 5,013,830; 5,149,797; 5,220,007; 5,256,775; 5,366,878; 5,403,711; 5,491,133; 5,565,350; 5,623,065; 5,652,355; 5,652,356; and 5,700,922, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference in its entirety.

The antisense compounds used in accordance with this invention may be conveniently and routinely made through the well-known technique of solid phase synthesis. Equipment for such synthesis is sold by several vendors including, for example, Applied Biosystems (Foster City, CA). Any other means for such synthesis known in the art may additionally or alternatively be employed. It is well known to use similar techniques to prepare oligonucleotides such as the phosphorothioates and alkylated derivatives.

The antisense compounds of the invention are synthesized in vitro and do not include antisense compositions of biological origin, or genetic vector constructs designed to direct the in vivo synthesis of antisense molecules.

The compounds of the invention may also be admixed, encapsulated, conjugated or otherwise associated with other molecules, molecule structures or mixtures of compounds, as for example, liposomes, receptor targeted molecules, oral, rectal, topical or other formulations, for assisting in uptake, distribution and/or absorption. Representative United States patents that teach the preparation of such uptake, distribution and/or absorption assisting

-20-

formulations include, but are not limited to, U.S.:

5,108,921; 5,354,844; 5,416,016; 5,459,127; 5,521,291;  
5,543,158; 5,547,932; 5,583,020; 5,591,721; 4,426,330;  
4,534,899; 5,013,556; 5,108,921; 5,213,804; 5,227,170;  
5 5,264,221; 5,356,633; 5,395,619; 5,416,016; 5,417,978;  
5,462,854; 5,469,854; 5,512,295; 5,527,528; 5,534,259;  
5,543,152; 5,556,948; 5,580,575; and 5,595,756, each of  
which is herein incorporated by reference.

The antisense compounds of the invention encompass  
10 any pharmaceutically acceptable salts, esters, or salts of  
such esters, or any other compound which, upon  
administration to an animal including a human, is capable  
of providing (directly or indirectly) the biologically  
active metabolite or residue thereof. Accordingly, for  
15 example, the disclosure is also drawn to prodrugs and  
pharmaceutically acceptable salts of the compounds of the  
invention, pharmaceutically acceptable salts of such  
prodrugs, and other bioequivalents.

The term "prodrug" indicates a therapeutic agent that  
20 is prepared in an inactive form that is converted to an  
active form (i.e., drug) within the body or cells thereof  
by the action of endogenous enzymes or other chemicals  
and/or conditions. In particular, prodrug versions of the  
~~original nucleotides of the invention are prepared as SATE~~  
25 [(S-acetyl-2-thioethyl) phosphate] derivatives according to  
the methods disclosed in WO 93/24510 to Gosselin et al.,  
published December 9, 1993 or in WO 94/26764 to Imbach et  
al.

The term "pharmaceutically acceptable salts" refers  
30 to physiologically and pharmaceutically acceptable salts of  
the compounds of the invention: i.e., salts that retain the  
desired biological activity of the parent compound and do  
not impart undesired toxicological effects thereto.

Pharmaceutically acceptable base addition salts are  
35 formed with metals or amines, such as alkali and alkaline

-21-

earth metals or organic amines. Examples of metals used as cations are sodium, potassium, magnesium, calcium, and the like. Examples of suitable amines are N,N'-dibenzylethylenediamine, chlorprocaine, choline, 5 diethanolamine, dicyclohexylamine, ethylenediamine, N-methylglucamine, and procaine (see, for example, Berge et al., "Pharmaceutical Salts," *J. of Pharma Sci.*, 1977, 66, 1-19). The base addition salts of said acidic compounds are prepared by contacting the free acid form with a 10 sufficient amount of the desired base to produce the salt in the conventional manner. The free acid form may be regenerated by contacting the salt form with an acid and isolating the free acid in the conventional manner. The free acid forms differ from their respective salt forms 15 somewhat in certain physical properties such as solubility in polar solvents, but otherwise the salts are equivalent to their respective free acid for purposes of the present invention. As used herein, a "pharmaceutical addition salt" includes a pharmaceutically acceptable salt of an 20 acid form of one of the components of the compositions of the invention. These include organic or inorganic acid salts of the amines. Preferred acid salts are the hydrochlorides, acetates, salicylates, nitrates and phosphates. Other suitable pharmaceutically acceptable 25 salts are well known to those skilled in the art and include basic salts of a variety of inorganic and organic acids, such as, for example, with inorganic acids, such as for example hydrochloric acid, hydrobromic acid, sulfuric acid or phosphoric acid; with organic carboxylic, sulfonic, 30 sulfo or phospho acids or N-substituted sulfamic acids, for example acetic acid, propionic acid, glycolic acid, succinic acid, maleic acid, hydroxymaleic acid, methylmaleic acid, fumaric acid, malic acid, tartaric acid, lactic acid, oxalic acid, gluconic acid, glucaric acid, 35 glucuronic acid, citric acid, benzoic acid, cinnamic acid,

-22-

mandelic acid, salicylic acid, 4-aminosalicylic acid, 2-phenoxybenzoic acid, 2-acetoxybenzoic acid, embonic acid, nicotinic acid or isonicotinic acid; and with amino acids, such as the 20 alpha-amino acids involved in the synthesis of proteins in nature, for example glutamic acid or aspartic acid, and also with phenylacetic acid, methanesulfonic acid, ethanesulfonic acid, 2-hydroxyethanesulfonic acid, ethane-1,2-disulfonic acid, benzenesulfonic acid, 4-methylbenzenesulfoic acid, naphthalene-2-sulfonic acid, naphthalene-1,5-disulfonic acid, 2- or 3-phosphoglycerate, glucose-6-phosphate, N-cyclohexylsulfamic acid (with the formation of cyclamates), or with other acid organic compounds, such as ascorbic acid. Pharmaceutically acceptable salts of compounds may also be prepared with a pharmaceutically acceptable cation. Suitable pharmaceutically acceptable cations are well known to those skilled in the art and include alkaline, alkaline earth, ammonium and quaternary ammonium cations. Carbonates or hydrogen carbonates are also possible.

For oligonucleotides, preferred examples of pharmaceutically acceptable salts include but are not limited to (a) salts formed with cations such as sodium, potassium, ammonium, magnesium, calcium, polyamines such as spermine and spermidine, etc.; (b) acid addition salts formed with inorganic acids, for example hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, nitric acid and the like; (c) salts formed with organic acids such as, for example, acetic acid, oxalic acid, tartaric acid, succinic acid, maleic acid, fumaric acid, gluconic acid, citric acid, malic acid, ascorbic acid, benzoic acid, tannic acid, palmitic acid, alginic acid, polyglutamic acid, naphthalenesulfonic acid, methanesulfonic acid, p-toluenesulfonic acid, naphthalenedisulfonic acid, polygalacturonic acid, and the like; and (d) salts formed



-23-

from elemental anions such as chlorine, bromine, and iodine.

The antisense compounds of the present invention can be utilized for diagnostics, therapeutics, prophylaxis and as research reagents and kits. For therapeutics, an animal, preferably a human, suspected of having a disease or disorder which can be treated by modulating the expression of PI3 kinase p110 beta is treated by administering antisense compounds in accordance with this invention. The compounds of the invention can be utilized in pharmaceutical compositions by adding an effective amount of an antisense compound to a suitable pharmaceutically acceptable diluent or carrier. Use of the antisense compounds and methods of the invention may also be useful prophylactically, e.g., to prevent or delay infection, inflammation or tumor formation, for example.

The antisense compounds of the invention are useful for research and diagnostics, because these compounds hybridize to nucleic acids encoding PI3 kinase p110 beta, enabling sandwich and other assays to easily be constructed to exploit this fact. Hybridization of the antisense oligonucleotides of the invention with a nucleic acid encoding PI3 kinase p110 beta can be detected by means known in the art. Such means may include conjugation of an enzyme to the oligonucleotide, radiolabelling of the oligonucleotide or any other suitable detection means. Kits using such detection means for detecting the level of PI3 kinase p110 beta in a sample may also be prepared.

The present invention also includes pharmaceutical compositions and formulations which include the antisense compounds of the invention. The pharmaceutical compositions of the present invention may be administered in a number of ways depending upon whether local or systemic treatment is desired and upon the area to be treated. Administration may be topical (including

-24-

ophthalmic and to mucous membranes including vaginal and rectal delivery), pulmonary, e.g., by inhalation or insufflation of powders or aerosols, including by nebulizer; intratracheal, intranasal, epidermal and transdermal), oral or parenteral. Parenteral administration includes intravenous, intraarterial, subcutaneous, intraperitoneal or intramuscular injection or infusion; or intracranial, e.g., intrathecal or intraventricular, administration. Oligonucleotides with at least one 2'-O-methoxyethyl modification are believed to be particularly useful for oral administration.

Pharmaceutical compositions and formulations for topical administration may include transdermal patches, ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable. Coated condoms, gloves and the like may also be useful.

Compositions and formulations for oral administration include powders or granules, suspensions or solutions in water or non-aqueous media, capsules, sachets or tablets. Thickeners, flavoring agents, diluents, emulsifiers, dispersing aids or binders may be desirable.

---

Compositions and formulations for parenteral, intrathecal or intraventricular administration may include sterile aqueous solutions which may also contain buffers, diluents and other suitable additives such as, but not limited to, penetration enhancers, carrier compounds and other pharmaceutically acceptable carriers or excipients.

Pharmaceutical compositions of the present invention include, but are not limited to, solutions, emulsions, and liposome-containing formulations. These compositions may be generated from a variety of components that include, but are not limited to, preformed liquids, self-emulsifying solids and self-emulsifying semisolids.

-25-

The pharmaceutical formulations of the present invention, which may conveniently be presented in unit dosage form, may be prepared according to conventional techniques well known in the pharmaceutical industry. Such techniques include the step of bringing into association the active ingredients with the pharmaceutical carrier(s) or excipient(s). In general the formulations are prepared by uniformly and intimately bringing into association the active ingredients with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product

The compositions of the present invention may be formulated into any of many possible dosage forms such as, but not limited to, tablets, capsules, liquid syrups, soft gels, suppositories, and enemas. The compositions of the present invention may also be formulated as suspensions in aqueous, non-aqueous or mixed media. Aqueous suspensions may further contain substances which increase the viscosity of the suspension including, for example, sodium carboxymethylcellulose, sorbitol and/or dextran. The suspension may also contain stabilizers.

In one embodiment of the present invention the pharmaceutical compositions may be formulated and used as foams. Pharmaceutical foams include formulations such as, but not limited to, emulsions, microemulsions, creams, jellies and liposomes. While basically similar in nature these formulations vary in the components and the consistency of the final product. The preparation of such compositions and formulations is generally known to those skilled in the pharmaceutical and formulation arts and may be applied to the formulation of the compositions of the present invention.

#### Emulsions

The compositions of the present invention may be prepared and formulated as emulsions. Emulsions are

-26-

typically heterogenous systems of one liquid dispersed in another in the form of droplets usually exceeding 0.1  $\mu\text{m}$  in diameter. (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199; Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., Volume 1, p. 245; Block in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 2, p. 335; Higuchi et al., in *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, PA, 1985, p. 301). Emulsions are often biphasic systems comprising of two immiscible liquid phases intimately mixed and dispersed with each other. In general, emulsions may be either water-in-oil (w/o) or of the oil-in-water (o/w) variety. When an aqueous phase is finely divided into and dispersed as minute droplets into a bulk oily phase the resulting composition is called a water-in-oil (w/o) emulsion. Alternatively, when an oily phase is finely divided into and dispersed as minute droplets into a bulk aqueous phase the resulting composition is called an oil-in-water (o/w) emulsion. Emulsions may contain additional components in addition to the dispersed phases and the active drug which may be present as a solution in either the aqueous phase, oily phase or itself as a separate phase. Pharmaceutical excipients such as emulsifiers, stabilizers, dyes, and anti-oxidants may also be present in emulsions as needed. Pharmaceutical emulsions may also be multiple emulsions that are comprised of more than two phases such as, for example, in the case of oil-in-water-in-oil (o/w/o) and water-in-oil-in-water (w/o/w) emulsions. Such complex formulations often provide certain advantages that simple binary emulsions do not. Multiple emulsions in which

-27-

individual oil droplets of an o/w emulsion enclose small water droplets constitute a w/o/w emulsion. Likewise a system of oil droplets enclosed in globules of water stabilized in an oily continuous provides an o/w/o  
5 emulsion.

Emulsions are characterized by little or no thermodynamic stability. Often, the dispersed or discontinuous phase of the emulsion is well dispersed into the external or continuous phase and maintained in this  
10 form through the means of emulsifiers or the viscosity of the formulation. Either of the phases of the emulsion may be a semisolid or a solid, as is the case of emulsion-style ointment bases and creams. Other means of stabilizing emulsions entail the use of emulsifiers that may be  
15 incorporated into either phase of the emulsion. Emulsifiers may broadly be classified into four categories: synthetic surfactants, naturally occurring emulsifiers, absorption bases, and finely dispersed solids (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker  
20 (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199).

Synthetic surfactants, also known as surface active agents, have found wide applicability in the formulation of emulsions and have been reviewed in the literature (Rieger,  
25 in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 285; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), Marcel Dekker, Inc., New York, N.Y., 1988, volume 1, p. 199). Surfactants are  
30 typically amphiphilic and comprise a hydrophilic and a hydrophobic portion. The ratio of the hydrophilic to the hydrophobic nature of the surfactant has been termed the hydrophile/lipophile balance (HLB) and is a valuable tool in categorizing and selecting surfactants in the  
35 preparation of formulations. Surfactants may be classified

-28-

into different classes based on the nature of the hydrophilic group: nonionic, anionic, cationic and amphoteric (Rieger, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 285).

Naturally occurring emulsifiers used in emulsion formulations include lanolin, beeswax, phosphatides, lecithin and acacia. Absorption bases possess hydrophilic properties such that they can soak up water to form w/o emulsions yet retain their semisolid consistencies, such as anhydrous lanolin and hydrophilic petrolatum. Finely divided solids have also been used as good emulsifiers especially in combination with surfactants and in viscous preparations. These include polar inorganic solids, such as heavy metal hydroxides, nonswelling clays such as bentonite, attapulgite, hectorite, kaolin, montmorillonite, colloidal aluminum silicate and colloidal magnesium aluminum silicate, pigments and nonpolar solids such as carbon or glyceryl tristearate.

A large variety of non-emulsifying materials are also included in emulsion formulations and contribute to the properties of emulsions. These include fats, oils, waxes, fatty acids, fatty alcohols, fatty esters, humectants, hydrophilic colloids, preservatives and antioxidants

(Block, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 335; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199).

Hydrophilic colloids or hydrocolloids include naturally occurring gums and synthetic polymers such as polysaccharides (for example, acacia, agar, alginic acid, carrageenan, guar gum, karaya gum, and tragacanth), cellulose derivatives (for example, carboxymethylcellulose and carboxypropylcellulose), and synthetic polymers (for

-29-

example, carbomers, cellulose ethers, and carboxyvinyl polymers). These disperse or swell in water to form colloidal solutions that stabilize emulsions by forming strong interfacial films around the dispersed-phase droplets and by increasing the viscosity of the external phase.

Since emulsions often contain a number of ingredients such as carbohydrates, proteins, sterols and phosphatides that may readily support the growth of microbes, these formulations often incorporate preservatives. Commonly used preservatives included in emulsion formulations include methyl paraben, propyl paraben, quaternary ammonium salts, benzalkonium chloride, esters of p-hydroxybenzoic acid, and boric acid. Antioxidants are also commonly added to emulsion formulations to prevent deterioration of the formulation. Antioxidants used may be free radical scavengers such as tocopherols, alkyl gallates, butylated hydroxyanisole, butylated hydroxytoluene, or reducing agents such as ascorbic acid and sodium metabisulfite, and antioxidant synergists such as citric acid, tartaric acid, and lecithin.

The application of emulsion formulations via dermatological, oral and parenteral routes and methods for their manufacture have been reviewed in the literature (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199). Emulsion formulations for oral delivery have been very widely used because of reasons of ease of formulation, efficacy from an absorption and bioavailability standpoint. (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199). Mineral-oil base laxatives, oil-

-30-

soluble vitamins and high fat nutritive preparations are among the materials that have commonly been administered orally as o/w emulsions.

In one embodiment of the present invention, the compositions of oligonucleotides and nucleic acids are formulated as microemulsions. A microemulsion may be defined as a system of water, oil and amphiphile which is a single optically isotropic and thermodynamically stable liquid solution (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245). Typically microemulsions are systems that are prepared by first dispersing an oil in an aqueous surfactant solution and then adding a sufficient amount of a fourth component, generally an intermediate chain-length alcohol to form a transparent system. Therefore, microemulsions have also been described as thermodynamically stable, isotropically clear dispersions of two immiscible liquids that are stabilized by interfacial films of surface-active molecules (Leung and Shah, in: *Controlled Release of Drugs: Polymers and Aggregate Systems*, Rosoff, M., Ed., 1989, VCH Publishers, New York, pages 185-215). Microemulsions commonly are prepared via a combination of three to five components that include oil, water, surfactant, cosurfactant and electrolyte. Whether the microemulsion is of the water-in-oil (w/o) or an oil-in-water (o/w) type is dependent on the properties of the oil and surfactant used and on the structure and geometric packing of the polar heads and hydrocarbon tails of the surfactant molecules (Schott, in *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, PA, 1985, p. 271).

The phenomenological approach utilizing phase diagrams has been extensively studied and has yielded a comprehensive knowledge, to one skilled in the art, of how



-31-

to formulate microemulsions (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245; Block, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 335). Compared to conventional emulsions, microemulsions offer the advantage of solubilizing water-insoluble drugs in a formulation of thermodynamically stable droplets that are formed spontaneously.

Surfactants used in the preparation of microemulsions include, but are not limited to, ionic surfactants, non-ionic surfactants, Brij 96, polyoxyethylene oleyl ethers, polyglycerol fatty acid esters, tetraglycerol monolaurate (ML310), tetraglycerol monooleate (MO310), hexaglycerol monooleate (PO310), hexaglycerol pentaoleate (PO500), decaglycerol monocaprate (MCA750), decaglycerol monooleate (MO750), decaglycerol sequioleate (SO750), decaglycerol decaoleate (DAO750), alone or in combination with cosurfactants. The cosurfactant, usually a short-chain alcohol such as ethanol, 1-propanol, and 1-butanol, serves to increase the interfacial fluidity by penetrating into the surfactant film and consequently creating a disordered film because of the void space generated among surfactant molecules. Microemulsions may, however, be prepared without the use of cosurfactants and alcohol-free self-emulsifying microemulsion systems are known in the art. The aqueous phase may typically be, but is not limited to, water, an aqueous solution of the drug, glycerol, PEG300, PEG400, polyglycerols, propylene glycols, and derivatives of ethylene glycol. The oil phase may include, but is not limited to, materials such as Captex 300, Captex 355, Capmul MCM, fatty acid esters, medium chain (C8-C12) mono, di, and tri-glycerides, polyoxyethylated glyceryl fatty acid esters, fatty alcohols, polyglycolized glycerides,

-32-

saturated polyglycolized C8-C10 glycerides, vegetable oils and silicone oil.

Microemulsions are particularly of interest from the standpoint of drug solubilization and the enhanced  
5 absorption of drugs. Lipid based microemulsions (both o/w and w/o) have been proposed to enhance the oral bioavailability of drugs, including peptides (Constantinides et al., *Pharmaceutical Research*, 1994, 11, 1385-1390; Ritschel, *Meth. Find. Exp. Clin. Pharmacol.*,  
10 1993, 13, 205). Microemulsions afford advantages of improved drug solubilization, protection of drug from enzymatic hydrolysis, possible enhancement of drug absorption due to surfactant-induced alterations in membrane fluidity and permeability, ease of preparation,  
15 ease of oral administration over solid dosage forms, improved clinical potency, and decreased toxicity (Constantinides et al., *Pharmaceutical Research*, 1994, 11, 1385; Ho et al., *J. Pharm. Sci.*, 1996, 85, 138-143). Often microemulsions may form spontaneously when their components  
20 are brought together at ambient temperature. This may be particularly advantageous when formulating thermolabile drugs, peptides or oligonucleotides. Microemulsions have also been effective in the transdermal delivery of active components in both cosmetic and pharmaceutical  
25 applications. It is expected that the microemulsion compositions and formulations of the present invention will facilitate the increased systemic absorption of oligonucleotides and nucleic acids from the gastrointestinal tract, as well as improve the local  
30 cellular uptake of oligonucleotides and nucleic acids within the gastrointestinal tract, vagina, buccal cavity and other areas of administration.

Microemulsions of the present invention may also contain additional components and additives such as

-33-

sorbitan monostearate (Grill 3), Labrasol, and penetration enhancers to improve the properties of the formulation and to enhance the absorption of the oligonucleotides and nucleic acids of the present invention. Penetration enhancers used in the microemulsions of the present invention may be classified as belonging to one of five broad categories - surfactants, fatty acids, bile salts, chelating agents, and non-chelating non-surfactants (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p. 92). Each of these classes has been discussed above.

#### Liposomes

There are many organized surfactant structures besides microemulsions that have been studied and used for the formulation of drugs. These include monolayers, micelles, bilayers and vesicles. Vesicles, such as liposomes, have attracted great interest because of their specificity and the duration of action they offer from the standpoint of drug delivery. As used in the present invention, the term "liposome" means a vesicle composed of amphiphilic lipids arranged in a spherical bilayer or bilayers.

Liposomes are unilamellar or multilamellar vesicles which have a membrane formed from a lipophilic material and an aqueous interior. The aqueous portion contains the composition to be delivered. Cationic liposomes possess the advantage of being able to fuse to the cell wall. Non-cationic liposomes, although not able to fuse as efficiently with the cell wall, are taken up by macrophages *in vivo*.

In order to cross intact mammalian skin, lipid vesicles must pass through a series of fine pores, each with a diameter less than 50 nm, under the influence of a suitable transdermal gradient. Therefore, it is desirable

-34-

to use a liposome which is highly deformable and able to pass through such fine pores.

Further advantages of liposomes include; liposomes obtained from natural phospholipids are biocompatible and biodegradable; liposomes can incorporate a wide range of water and lipid soluble drugs; liposomes can protect encapsulated drugs in their internal compartments from metabolism and degradation (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245). Important considerations in the preparation of liposome formulations are the lipid surface charge, vesicle size and the aqueous volume of the liposomes.

Liposomes are useful for the transfer and delivery of active ingredients to the site of action. Because the liposomal membrane is structurally similar to biological membranes, when liposomes are applied to a tissue, the liposomes start to merge with the cellular membranes. As the merging of the liposome and cell progresses, the liposomal contents are emptied into the cell where the active agent may act.

Liposomal formulations have been the focus of extensive investigation as the mode of delivery for many drugs. There is growing evidence that for topical administration, liposomes present several advantages over other formulations. Such advantages include reduced side-effects related to high systemic absorption of the administered drug, increased accumulation of the administered drug at the desired target, and the ability to administer a wide variety of drugs, both hydrophilic and hydrophobic, into the skin.

Several reports have detailed the ability of liposomes to deliver agents including high-molecular weight DNA into the skin. Compounds including analgesics, antibodies, hormones and high-molecular weight DNAs have

-35-

been administered to the skin. The majority of applications resulted in the targeting of the upper epidermis.

Liposomes fall into two broad classes. Cationic liposomes are positively charged liposomes which interact with the negatively charged DNA molecules to form a stable complex. The positively charged DNA/liposome complex binds to the negatively charged cell surface and is internalized in an endosome. Due to the acidic pH within the endosome, the liposomes are ruptured, releasing their contents into the cell cytoplasm (Wang et al., *Biochem. Biophys. Res. Commun.*, 1987, 147, 980-985).

Liposomes which are pH-sensitive or negatively-charged, entrap DNA rather than complex with it. Since both the DNA and the lipid are similarly charged, repulsion rather than complex formation occurs. Nevertheless, some DNA is entrapped within the aqueous interior of these liposomes. pH-sensitive liposomes have been used to deliver DNA encoding the thymidine kinase gene to cell monolayers in culture. Expression of the exogenous gene was detected in the target cells (Zhou et al., *Journal of Controlled Release*, 1992, 19, 269-274).

One major type of liposomal composition includes phospholipids other than naturally-derived phosphatidylcholine. Neutral liposome compositions, for example, can be formed from dimyristoyl phosphatidylcholine (DMPC) or dipalmitoyl phosphatidylcholine (DPPC). Anionic liposome compositions generally are formed from dimyristoyl phosphatidylglycerol, while anionic fusogenic liposomes are formed primarily from dioleoyl phosphatidylethanolamine (DOPE). Another type of liposomal composition is formed from phosphatidylcholine (PC) such as, for example, soybean PC, and egg PC. Another type is formed from mixtures of phospholipid and/or phosphatidylcholine and/or cholesterol.

-36-

Several studies have assessed the topical delivery of liposomal drug formulations to the skin. Application of liposomes containing interferon to guinea pig skin resulted in a reduction of skin herpes sores while delivery of  
5 interferon via other means (e.g. as a solution or as an emulsion) were ineffective (Weiner et al., *Journal of Drug Targeting*, 1992, 2, 405-410). Further, an additional study tested the efficacy of interferon administered as part of a liposomal formulation to the administration of interferon  
10 using an aqueous system, and concluded that the liposomal formulation was superior to aqueous administration (du Plessis et al., *Antiviral Research*, 1992, 18, 259-265).

Non-ionic liposomal systems have also been examined to determine their utility in the delivery of drugs to the  
15 skin, in particular systems comprising non-ionic surfactant and cholesterol. Non-ionic liposomal formulations comprising Novasome™ I (glyceryl dilaurate/cholesterol/polyoxyethylene-10-stearyl ether) and Novasome™ II (glyceryl distearate/  
20 cholesterol/polyoxyethylene-10-stearyl ether) were used to deliver cyclosporin-A into the dermis of mouse skin. Results indicated that such non-ionic liposomal systems were effective in facilitating the deposition of  
cyclosporin-A into different layers of the skin (Hu et al.  
25 *S.T.P. Pharma. Sci.*, 1994, 4, 6, 466).

Liposomes also include "sterically stabilized" liposomes, a term which, as used herein, refers to liposomes comprising one or more specialized lipids that, when incorporated into liposomes, result in enhanced  
30 circulation lifetimes relative to liposomes lacking such specialized lipids. Examples of sterically stabilized liposomes are those in which part of the vesicle-forming lipid portion of the liposome (A) comprises one or more glycolipids, such as monosialoganglioside G<sub>M1</sub>, or (B) is  
35 derivatized with one or more hydrophilic polymers, such as

-37-

a polyethylene glycol (PEG) moiety. While not wishing to be bound by any particular theory, it is thought in the art that, at least for sterically stabilized liposomes containing gangliosides, sphingomyelin, or PEG-derivatized lipids, the enhanced circulation half-life of these sterically stabilized liposomes derives from a reduced uptake into cells of the reticuloendothelial system (RES) (Allen et al., *FEBS Letters*, 1987, 223, 42; Wu et al., *Cancer Research*, 1993, 53, 3765). Various liposomes comprising one or more glycolipids are known in the art. Papahadjopoulos et al. (*Ann. N.Y. Acad. Sci.*, 1987, 507, 64) reported the ability of monosialoganglioside  $G_{M1}$ , galactocerebroside sulfate and phosphatidylinositol to improve blood half-lives of liposomes. These findings were expounded upon by Gabizon et al. (*Proc. Natl. Acad. Sci. U.S.A.*, 1988, 85, 6949). U.S. Patent No. 4,837,028 and WO 88/04924, both to Allen et al., disclose liposomes comprising (1) sphingomyelin and (2) the ganglioside  $G_{M1}$  or a galactocerebroside sulfate ester. U.S. Patent No. 5,543,152 (Webb et al.) discloses liposomes comprising sphingomyelin. Liposomes comprising 1,2-sn-dimyristoylphosphatidylcholine are disclosed in WO 97/13499 (Lim et al.).

Many liposomes comprising lipids derivatized with one or more hydrophilic polymers, and methods of preparation thereof, are known in the art. Sunamoto et al. (*Bull. Chem. Soc. Jpn.*, 1980, 53, 2778) described liposomes comprising a nonionic detergent, 2C<sub>12</sub>15G, that contains a PEG moiety. Illum et al. (*FEBS Lett.*, 1984, 167, 79) noted that hydrophilic coating of polystyrene particles with polymeric glycols results in significantly enhanced blood half-lives. Synthetic phospholipids modified by the attachment of carboxylic groups of polyalkylene glycols (e.g., PEG) are described by Sears (U.S. Patent Nos.

-38-

4,426,330 and 4,534,899). Klibanov et al. (*FEBS Lett.*, 1990, 268, 235) described experiments demonstrating that liposomes comprising phosphatidylethanolamine (PE) derivatized with PEG or PEG stearate have significant increases in blood circulation half-lives. Blume et al. (*Biochimica et Biophysica Acta*, 1990, 1029, 91) extended such observations to other PEG-derivatized phospholipids, e.g., DSPE-PEG, formed from the combination of distearoylphosphatidylethanolamine (DSPE) and PEG.

10 Liposomes having covalently bound PEG moieties on their external surface are described in European Patent No. EP 0 445 131 B1 and WO 90/04384 to Fisher. Liposome compositions containing 1-20 mole percent of PE derivatized with PEG, and methods of use thereof, are described by

15 Woodle et al. (U.S. Patent Nos. 5,013,556 and 5,356,633) and Martin et al. (U.S. Patent No. 5,213,804 and European Patent No. EP 0 496 813 B1). Liposomes comprising a number of other lipid-polymer conjugates are disclosed in WO 91/05545 and U.S. Patent No. 5,225,212 (both to Martin et

20 al.) and in WO 94/20073 (Zalipsky et al.) Liposomes comprising PEG-modified ceramide lipids are described in WO 96/10391 (Choi et al.). U.S. Patent Nos. 5,540,935 (Miyazaki et al.) and 5,556,948 (Tagawa et al.) describe

PEG-containing liposomes that can be further derivatized

25 with functional moieties on their surfaces.

A limited number of liposomes comprising nucleic acids are known in the art. WO 96/40062 to Thierry et al. discloses methods for encapsulating high molecular weight nucleic acids in liposomes. U.S. Patent No. 5,264,221 to

30 Tagawa et al. discloses protein-bonded liposomes and asserts that the contents of such liposomes may include an antisense RNA. U.S. Patent No. 5,665,710 to Rahman et al. describes certain methods of encapsulating oligodeoxynucleotides in liposomes. WO 97/04787 to Love et



-39-

al. discloses liposomes comprising antisense oligonucleotides targeted to the raf gene.

Transfersomes are yet another type of liposomes, and are highly deformable lipid aggregates which are attractive candidates for drug delivery vehicles. Transfersomes may be described as lipid droplets which are so highly deformable that they are easily able to penetrate through pores which are smaller than the droplet. Transfersomes are adaptable to the environment in which they are used, e.g. they are self-optimizing (adaptive to the shape of pores in the skin), self-repairing, frequently reach their targets without fragmenting, and often self-loading. To make transfersomes it is possible to add surface edge-activators, usually surfactants, to a standard liposomal composition. Transfersomes have been used to deliver serum albumin to the skin. The transfersome-mediated delivery of serum albumin has been shown to be as effective as subcutaneous injection of a solution containing serum albumin.

Surfactants find wide application in formulations such as emulsions (including microemulsions) and liposomes. The most common way of classifying and ranking the properties of the many different types of surfactants, both natural and synthetic, is by the use of the hydrophile/lipophile balance (HLB). The nature of the hydrophilic group (also known as the "head") provides the most useful means for categorizing the different surfactants used in formulations (Rieger, in *Pharmaceutical Dosage Forms*, Marcel Dekker, Inc., New York, NY, 1988, p. 285).

If the surfactant molecule is not ionized, it is classified as a nonionic surfactant. Nonionic surfactants find wide application in pharmaceutical and cosmetic products and are usable over a wide range of pH values. In general their HLB values range from 2 to about 18 depending

on their structure. Nonionic surfactants include nonionic esters such as ethylene glycol esters, propylene glycol esters, glyceryl esters, polyglyceryl esters, sorbitan esters, sucrose esters, and ethoxylated esters. Nonionic  
5 alkanolamides and ethers such as fatty alcohol ethoxylates, propoxylated alcohols, and ethoxylated/propoxylated block polymers are also included in this class. The polyoxyethylene surfactants are the most popular members of the nonionic surfactant class.

10 If the surfactant molecule carries a negative charge when it is dissolved or dispersed in water, the surfactant is classified as anionic. Anionic surfactants include carboxylates such as soaps, acyl lactylates, acyl amides of amino acids, esters of sulfuric acid such as alkyl sulfates  
15 and ethoxylated alkyl sulfates, sulfonates such as alkyl benzene sulfonates, acyl isethionates, acyl taurates and sulfosuccinates, and phosphates. The most important members of the anionic surfactant class are the alkyl sulfates and the soaps.

20 If the surfactant molecule carries a positive charge when it is dissolved or dispersed in water, the surfactant is classified as cationic. Cationic surfactants include quaternary ammonium salts and ethoxylated amines. The  

---

quaternary ammonium salts are the most used members of this  
25 class.

If the surfactant molecule has the ability to carry either a positive or negative charge, the surfactant is classified as amphoteric. Amphoteric surfactants include acrylic acid derivatives, substituted alkylamides, N-  
30 alkylbetaines and phosphatides.

The use of surfactants in drug products, formulations and in emulsions has been reviewed (Rieger, in *Pharmaceutical Dosage Forms*, Marcel Dekker, Inc., New York, NY, 1988, p. 285).

-41-

### Penetration Enhancers

In one embodiment, the present invention employs various penetration enhancers to effect the efficient delivery of nucleic acids, particularly oligonucleotides, to the skin of animals. Most drugs are present in solution in both ionized and nonionized forms. However, usually only lipid soluble or lipophilic drugs readily cross cell membranes. It has been discovered that even non-lipophilic drugs may cross cell membranes if the membrane to be crossed is treated with a penetration enhancer. In addition to aiding the diffusion of non-lipophilic drugs across cell membranes, penetration enhancers also enhance the permeability of lipophilic drugs.

Penetration enhancers may be classified as belonging to one of five broad categories, i.e., surfactants, fatty acids, bile salts, chelating agents, and non-chelating non-surfactants (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p.92). Each of the above mentioned classes of penetration enhancers are described below in greater detail.

Surfactants: In connection with the present invention, surfactants (or "surface-active agents") are chemical entities which, when dissolved in an aqueous solution, reduce the surface tension of the solution or the interfacial tension between the aqueous solution and another liquid, with the result that absorption of oligonucleotides through the mucosa is enhanced. In addition to bile salts and fatty acids, these penetration enhancers include, for example, sodium lauryl sulfate, polyoxyethylene-9-lauryl ether and polyoxyethylene-20-cetyl ether) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p.92); and perfluorochemical emulsions, such as FC-43. Takahashi et al., *J. Pharm. Pharmacol.*, 1988, 40, 252).

-42-

Fatty acids: Various fatty acids and their derivatives which act as penetration enhancers include, for example, oleic acid, lauric acid, capric acid (n-decanoic acid), myristic acid, palmitic acid, stearic acid, linoleic acid, linolenic acid, dicaprate, tricaprate, monoolein (1-monooleoyl-rac-glycerol), dilaurin, caprylic acid, arachidonic acid, glycerol 1-monocaprate, 1-dodecylazacycloheptan-2-one, acylcarnitines, acylcholines, C<sub>1-10</sub> alkyl esters thereof (e.g., methyl, isopropyl and t-butyl), and mono- and di-glycerides thereof (i.e., oleate, laurate, caprate, myristate, palmitate, stearate, linoleate, etc.) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p.92; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33; El Hariri et al., *J. Pharm. Pharmacol.*, 1992, 44, 651-654).

Bile salts: The physiological role of bile includes the facilitation of dispersion and absorption of lipids and fat-soluble vitamins (Brunton, Chapter 38 in: Goodman & Gilman's *The Pharmacological Basis of Therapeutics*, 9th Ed., Hardman et al. Eds., McGraw-Hill, New York, 1996, pp. 934-935). Various natural bile salts, and their synthetic derivatives, act as penetration enhancers. Thus the term "bile salts" includes any of the naturally occurring components of bile as well as any of their synthetic derivatives. The bile salts of the invention include, for example, cholic acid (or its pharmaceutically acceptable sodium salt, sodium cholate), dehydrocholic acid (sodium dehydrocholate), deoxycholic acid (sodium deoxycholate), glucolic acid (sodium glucolate), glycholic acid (sodium glycocholate), glycodeoxycholic acid (sodium glycodeoxycholate), taurocholic acid (sodium taurocholate), taurodeoxycholic acid (sodium taurodeoxycholate), chenodeoxycholic acid (sodium chenodeoxycholate),

-43-

- ursodeoxycholic acid (UDCA), sodium tauro-24,25-dihydrofusidate (STDHF), sodium glycodihydrofusidate and polyoxyethylene-9-lauryl ether (POE) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, page 92;
- 5 Swinyard, Chapter 39 In: *Remington's Pharmaceutical Sciences*, 18th Ed., Gennaro, ed., Mack Publishing Co., Easton, PA, 1990, pages 782-783; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33; Yamamoto et al., *J. Pharm. Exp. Ther.*, 1992, 263, 25;
- 10 Yamashita et al., *J. Pharm. Sci.*, 1990, 79, 579-583).

Chelating Agents: Chelating agents, as used in connection with the present invention, can be defined as compounds that remove metallic ions from solution by forming complexes therewith, with the result that

15 absorption of oligonucleotides through the mucosa is enhanced. With regards to their use as penetration enhancers in the present invention, chelating agents have the added advantage of also serving as DNase inhibitors, as most characterized DNA nucleases require a divalent metal

20 ion for catalysis and are thus inhibited by chelating agents (Jarrett, *J. Chromatogr.*, 1993, 618, 315-339). Chelating agents of the invention include but are not limited to disodium ethylenediaminetetraacetate (EDTA), citric acid, salicylates (e.g., sodium salicylate, 5-

25 methoxysalicylate and homovanilate), *N*-acyl derivatives of collagen, laureth-9 and *N*-amino acyl derivatives of beta-diketones (enamines) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, page 92; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990,

30 7, 1-33; Buur et al., *J. Control Rel.*, 1990, 14, 43-51).

Non-chelating non-surfactants: As used herein, non-chelating non-surfactant penetration enhancing compounds can be defined as compounds that demonstrate insignificant activity as chelating agents or as surfactants but that

-44-

nonetheless enhance absorption of oligonucleotides through the alimentary mucosa (Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33). This class of penetration enhancers include, for example, 5 unsaturated cyclic ureas, 1-alkyl- and 1-alkenylazacyclo-alkanone derivatives (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, page 92); and non-steroidal anti-inflammatory agents such as diclofenac sodium, indomethacin and phenylbutazone (Yamashita et al., 10 *J. Pharm. Pharmacol.*, 1987, 39, 621-626).

Agents that enhance uptake of oligonucleotides at the cellular level may also be added to the pharmaceutical and other compositions of the present invention. For example, cationic lipids, such as lipofectin (Junichi et al, U.S. 15 Patent No. 5,705,188), cationic glycerol derivatives, and polycationic molecules, such as polylysine (Lollo et al., PCT Application WO 97/30731), are also known to enhance the cellular uptake of oligonucleotides.

Other agents may be utilized to enhance the 20 penetration of the administered nucleic acids, including glycols such as ethylene glycol and propylene glycol, pyrrols such as 2-pyrrol, azones, and terpenes such as limonene and menthone.

---

#### Carriers

25 Certain compositions of the present invention also incorporate carrier compounds in the formulation. As used herein, "carrier compound" or "carrier" can refer to a nucleic acid, or analog thereof, which is inert (i.e., does not possess biological activity per se) but is recognized 30 as a nucleic acid by *in vivo* processes that reduce the bioavailability of a nucleic acid having biological activity by, for example, degrading the biologically active nucleic acid or promoting its removal from circulation. The coadministration of a nucleic acid and a carrier

-45-

compound, typically with an excess of the latter substance, can result in a substantial reduction of the amount of nucleic acid recovered in the liver, kidney or other extracirculatory reservoirs, presumably due to competition  
5 between the carrier compound and the nucleic acid for a common receptor. For example, the recovery of a partially phosphorothioate oligonucleotide in hepatic tissue can be reduced when it is coadministered with polyinosinic acid, dextran sulfate, polycytidic acid or 4-acetamido-  
10 4'-isothiocyano-stilbene-2,2'-disulfonic acid (Miyao et al., *Antisense Res. Dev.*, 1995, 5, 115-121; Takakura et al., *Antisense & Nucl. Acid Drug Dev.*, 1996, 6, 177-183).

#### Excipients

In contrast to a carrier compound, a "pharmaceutical  
15 carrier" or "excipient" is a pharmaceutically acceptable solvent, suspending agent or any other pharmacologically inert vehicle for delivering one or more nucleic acids to an animal. The excipient may be liquid or solid and is selected, with the planned manner of administration in  
20 mind, so as to provide for the desired bulk, consistency, etc., when combined with a nucleic acid and the other components of a given pharmaceutical composition. Typical pharmaceutical carriers include, but are not limited to, binding agents (e.g., pregelatinized maize starch,  
25 polyvinylpyrrolidone or hydroxypropyl methylcellulose, etc.); fillers (e.g., lactose and other sugars, microcrystalline cellulose, pectin, gelatin, calcium sulfate, ethyl cellulose, polyacrylates or calcium hydrogen phosphate, etc.); lubricants (e.g., magnesium stearate,  
30 talc, silica, colloidal silicon dioxide, stearic acid, metallic stearates, hydrogenated vegetable oils, corn starch, polyethylene glycols, sodium benzoate, sodium acetate, etc.); disintegrants (e.g., starch, sodium starch

-46-

glycolate, etc.); and wetting agents (e.g., sodium lauryl sulphate, etc.).

Pharmaceutically acceptable organic or inorganic excipient suitable for non-parenteral administration which  
5 do not deleteriously react with nucleic acids can also be used to formulate the compositions of the present invention. Suitable pharmaceutically acceptable carriers include, but are not limited to, water, salt solutions, alcohols, polyethylene glycols, gelatin, lactose, amylose,  
10 magnesium stearate, talc, silicic acid, viscous paraffin, hydroxymethylcellulose, polyvinylpyrrolidone and the like.

Formulations for topical administration of nucleic acids may include sterile and non-sterile aqueous solutions, non-aqueous solutions in common solvents such as  
15 alcohols, or solutions of the nucleic acids in liquid or solid oil bases. The solutions may also contain buffers, diluents and other suitable additives. Pharmaceutically acceptable organic or inorganic excipients suitable for non-parenteral administration which do not deleteriously  
20 react with nucleic acids can be used.

Suitable pharmaceutically acceptable excipients include, but are not limited to, water, salt solutions, alcohol, polyethylene glycols, gelatin, lactose, amylose,  
~~magnesium stearate, talc, silicic acid, viscous paraffin,~~  
25 hydroxymethylcellulose, polyvinylpyrrolidone and the like.

#### Other Components

The compositions of the present invention may additionally contain other adjunct components conventionally found in pharmaceutical compositions, at  
30 their art-established usage levels. Thus, for example, the compositions may contain additional, compatible, pharmaceutically-active materials such as, for example, antipruritics, astringents, local anesthetics or anti-inflammatory agents, or may contain additional  
35 materials useful in physically formulating various dosage .



-47-

forms of the compositions of the present invention, such as dyes, flavoring agents, preservatives, antioxidants, opacifiers, thickening agents and stabilizers. However, such materials, when added, should not unduly interfere with the biological activities of the components of the compositions of the present invention. The formulations can be sterilized and, if desired, mixed with auxiliary agents, e.g., lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, colorings, flavorings and/or aromatic substances and the like which do not deleteriously interact with the nucleic acid(s) of the formulation.

Aqueous suspensions may contain substances which increase the viscosity of the suspension including, for example, sodium carboxymethylcellulose, sorbitol and/or dextran. The suspension may also contain stabilizers.

Certain embodiments of the invention provide pharmaceutical compositions containing (a) one or more antisense compounds and (b) one or more other chemotherapeutic agents which function by a non-antisense mechanism. Examples of such chemotherapeutic agents include, but are not limited to, anticancer drugs such as daunorubicin, dactinomycin, doxorubicin, bleomycin, mitomycin, nitrogen mustard, chlorambucil, melphalan, cyclophosphamide, 6-mercaptopurine, 6-thioguanine, cytarabine (CA), 5-fluorouracil (5-FU), floxuridine (5-FUdR), methotrexate (MTX), colchicine, vincristine, vinblastine, etoposide, teniposide, cisplatin and diethylstilbestrol (DES). See, generally, *The Merck Manual of Diagnosis and Therapy*, 15th Ed., Berkow et al., eds., 1987, Rahway, N.J., pages 1206-1228). Anti-inflammatory drugs, including but not limited to nonsteroidal anti-inflammatory drugs and corticosteroids, and antiviral drugs, including but not limited to ribivirin, vidarabine, acyclovir and ganciclovir, may also be combined in

-48-

compositions of the invention. See, generally, *The Merck Manual of Diagnosis and Therapy*, 15th Ed., Berkow et al., eds., 1987, Rahway, N.J., pages 2499-2506 and 46-49, respectively). Other non-antisense chemotherapeutic agents  
5 are also within the scope of this invention. Two or more combined compounds may be used together or sequentially.

In another related embodiment, compositions of the invention may contain one or more antisense compounds, particularly oligonucleotides, targeted to a first nucleic  
10 acid and one or more additional antisense compounds targeted to a second nucleic acid target. Numerous examples of antisense compounds are known in the art. Two or more combined compounds may be used together or sequentially.

The formulation of therapeutic compositions and their  
15 subsequent administration is believed to be within the skill of those in the art. Dosing is dependent on severity and responsiveness of the disease state to be treated, with the course of treatment lasting from several days to several months, or until a cure is effected or a diminution  
20 of the disease state is achieved. Optimal dosing schedules can be calculated from measurements of drug accumulation in the body of the patient. Persons of ordinary skill can easily determine optimum dosages, dosing methodologies and  
~~repetition rates. Optimum dosages may vary depending on~~  
25 the relative potency of individual oligonucleotides, and can generally be estimated based on  $EC_{50}$ s found to be effective in in vitro and in vivo animal models. In general, dosage is from 0.01 ug to 100 g per kg of body weight, and may be given once or more daily, weekly,  
30 monthly or yearly, or even once every 2 to 20 years. Persons of ordinary skill in the art can easily estimate repetition rates for dosing based on measured residence times and concentrations of the drug in bodily fluids or tissues. Following successful treatment, it may be  
35 desirable to have the patient undergo maintenance therapy

-49-

to prevent the recurrence of the disease state, wherein the oligonucleotide is administered in maintenance doses, ranging from 0.01 ug to 100 g per kg of body weight, once or more daily, to once every 20 years.

5 While the present invention has been described with specificity in accordance with certain of its preferred embodiments, the following examples serve only to illustrate the invention and are not intended to limit the same.

## 10 EXAMPLES

### Example 1

#### Nucleoside Phosphoramidites for Oligonucleotide Synthesis Deoxy and 2'-alkoxy amidites

2'-Deoxy and 2'-methoxy beta-cyanoethyldiisopropyl  
15 phosphoramidites were purchased from commercial sources (e.g. Chemgenes, Needham MA or Glen Research, Inc. Sterling VA). Other 2'-O-alkoxy substituted nucleoside amidites are prepared as described in U.S. Patent 5,506,351, herein incorporated by reference. For oligonucleotides  
20 synthesized using 2'-alkoxy amidites, the standard cycle for unmodified oligonucleotides was utilized, except the wait step after pulse delivery of tetrazole and base was increased to 360 seconds.

Oligonucleotides containing 5-methyl-2'-deoxycytidine  
25 (5-Me-C) nucleotides were synthesized according to published methods [Sanghvi, et. al., *Nucleic Acids Research*, 1993, 21, 3197-3203] using commercially available phosphoramidites (Glen Research, Sterling VA or ChemGenes, Needham MA).

### 30 2'-Fluoro amidites

#### 2'-Fluorodeoxyadenosine amidites

2'-fluoro oligonucleotides were synthesized as described previously [Kawasaki, et. al., *J. Med. Chem.*, 1993, 36, 831-841] and United States patent 5,670,633,

-50-

herein incorporated by reference. Briefly, the protected nucleoside N6-benzoyl-2'-deoxy-2'-fluoroadenosine was synthesized utilizing commercially available 9-beta-D-arabinofuranosyladenine as starting material and by  
5 modifying literature procedures whereby the 2'-alpha-fluoro atom is introduced by a  $S_N2$ -displacement of a 2'-beta-trityl group. Thus N6-benzoyl-9-beta-D-arabinofuranosyladenine was selectively protected in moderate yield as the 3',5'-ditetrahydropyranyl (THP) intermediate. Deprotection of  
10 the THP and N6-benzoyl groups was accomplished using standard methodologies and standard methods were used to obtain the 5'-dimethoxytrityl-(DMT) and 5'-DMT-3'-phosphoramidite intermediates.

#### 2'-Fluorodeoxyguanosine

15 The synthesis of 2'-deoxy-2'-fluoroguanosine was accomplished using tetraisopropylidisiloxanyl (TPDS) protected 9-beta-D-arabinofuranosylguanine as starting material, and conversion to the intermediate diisobutyryl-arabinofuranosylguanosine. Deprotection of the TPDS group  
20 was followed by protection of the hydroxyl group with THP to give diisobutyryl di-THP protected arabinofuranosylguanine. Selective O-deacylation and triflation was followed by treatment of the crude product  
with fluoride, then deprotection of the THP groups.

25 Standard methodologies were used to obtain the 5'-DMT- and 5'-DMT-3'-phosphoramidites.

#### 2'-Fluorouridine

Synthesis of 2'-deoxy-2'-fluorouridine was accomplished by the modification of a literature procedure  
30 in which 2,2'-anhydro-1-beta-D-arabinofuranosyluracil was treated with 70% hydrogen fluoride-pyridine. Standard procedures were used to obtain the 5'-DMT and 5'-DMT-3'-phosphoramidites.

-51-

**2'-Fluorodeoxycytidine**

2'-deoxy-2'-fluorocytidine was synthesized via amination of 2'-deoxy-2'-fluorouridine, followed by selective protection to give N4-benzoyl-2'-deoxy-2'-fluorocytidine. Standard procedures were used to obtain the 5'-DMT and 5'-DMT-3'phosphoramidites.

**2'-O-(2-Methoxyethyl) modified amidites**

2'-O-Methoxyethyl-substituted nucleoside amidites are prepared as follows, or alternatively, as per the methods of Martin, P., *Helvetica Chimica Acta*, 1995, 78, 486-504.

**2,2'-Anhydro[1-(beta-D-arabinofuranosyl)-5-methyluridine]**

5-Methyluridine (ribosylthymine, commercially available through Yamasa, Choshi, Japan) (72.0 g, 0.279 M), diphenylcarbonate (90.0 g, 0.420 M) and sodium bicarbonate (2.0 g, 0.024 M) were added to DMF (300 mL). The mixture was heated to reflux, with stirring, allowing the evolved carbon dioxide gas to be released in a controlled manner. After 1 hour, the slightly darkened solution was concentrated under reduced pressure. The resulting syrup was poured into diethylether (2.5 L), with stirring. The product formed a gum. The ether was decanted and the residue was dissolved in a minimum amount of methanol (ca. 400 mL). The solution was poured into fresh ether (2.5 L) to yield a stiff gum. The ether was decanted and the gum was dried in a vacuum oven (60°C at 1 mm Hg for 24 h) to give a solid that was crushed to a light tan powder (57 g, 85% crude yield). The NMR spectrum was consistent with the structure, contaminated with phenol as its sodium salt (ca. 5%). The material was used as is for further reactions (or it can be purified further by column chromatography using a gradient of methanol in ethyl acetate (10-25%) to give a white solid, mp 222-4°C).

-52-

**2'-O-Methoxyethyl-5-methyluridine**

2,2'-Anhydro-5-methyluridine (195 g, 0.81 M), tris(2-methoxyethyl)borate (231 g, 0.98 M) and 2-methoxyethanol (1.2 L) were added to a 2 L stainless steel pressure vessel and placed in a pre-heated oil bath at 160°C. After heating for 48 hours at 155-160°C, the vessel was opened and the solution evaporated to dryness and triturated with MeOH (200 mL). The residue was suspended in hot acetone (1 L). The insoluble salts were filtered, washed with acetone (150 mL) and the filtrate evaporated. The residue (280 g) was dissolved in CH<sub>3</sub>CN (600 mL) and evaporated. A silica gel column (3 kg) was packed in CH<sub>2</sub>Cl<sub>2</sub>/acetone/MeOH (20:5:3) containing 0.5% Et<sub>3</sub>NH. The residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (250 mL) and adsorbed onto silica (150 g) prior to loading onto the column. The product was eluted with the packing solvent to give 160 g (63%) of product. Additional material was obtained by reworking impure fractions.

**2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine**

2'-O-Methoxyethyl-5-methyluridine (160 g, 0.506 M) was co-evaporated with pyridine (250 mL) and the dried residue dissolved in pyridine (1.3 L). A first aliquot of dimethoxytrityl chloride (94.3 g, 0.278 M) was added and the mixture stirred at room temperature for one hour. A

---

second aliquot of dimethoxytrityl chloride (94.3 g, 0.278 M) was added and the reaction stirred for an additional one hour. Methanol (170 mL) was then added to stop the reaction. HPLC showed the presence of approximately 70% product. The solvent was evaporated and triturated with CH<sub>3</sub>CN (200 mL). The residue was dissolved in CHCl<sub>3</sub> (1.5 L) and extracted with 2x500 mL of saturated NaHCO<sub>3</sub> and 2x500 mL of saturated NaCl. The organic phase was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and evaporated. 275 g of residue was obtained. The residue was purified on a 3.5 kg silica gel column, packed and eluted with EtOAc/hexane/acetone (5:5:1) containing 0.5% Et<sub>3</sub>NH. The pure fractions were evaporated

WO 01/18023

-53-

to give 164 g of product. Approximately 20 g additional was obtained from the impure fractions to give a total yield of 183 g (57%).

5 3'-O-Acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine

2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine (106 g, 0.167 M), DMF/pyridine (750 mL of a 3:1 mixture prepared from 562 mL of DMF and 188 mL of pyridine) and acetic anhydride (24.38 mL, 0.258 M) were combined and stirred at room temperature for 24 hours. The reaction was monitored by TLC by first quenching the TLC sample with the addition of MeOH. Upon completion of the reaction, as judged by TLC, MeOH (50 mL) was added and the mixture evaporated at 35°C. The residue was dissolved in CHCl<sub>3</sub> (800 mL) and extracted with 2x200 mL of saturated sodium bicarbonate and 2x200 mL of saturated NaCl. The water layers were back extracted with 200 mL of CHCl<sub>3</sub>, (800 combined organics were dried with sodium sulfate and evaporated to give 122 g of residue (approx. 90% product).

15 mL) and extracted with 200 mL of CHCl<sub>3</sub>. The water layers were back extracted with 200 mL of CHCl<sub>3</sub>. The combined organics were dried with sodium sulfate and evaporated to give 122 g of residue (approx. 90% product).

20 The residue was purified on a 3.5 kg silica gel column and eluted using EtOAc/hexane(4:1). Pure product fractions were evaporated to yield 96 g (84%). An additional 1.5 g was recovered from later fractions.

3'-O-Acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyl-4-triazoleuridine

25 A first solution was prepared by dissolving 3'-O-acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine (96 g, 0.144 M) in CH<sub>3</sub>CN (700 mL) and set aside. Triethylamine (189 mL, 1.44 M) was added to a solution of triazole (90 g, 1.3 M) in CH<sub>3</sub>CN (1 L), cooled to -5°C and stirred for 0.5 h using an overhead stirrer. POCl<sub>3</sub> was added dropwise, over a 30 minute period, to the stirred solution maintained at 0-10°C, and the resulting mixture stirred for an additional 2 hours. The first solution was 35 added dropwise, over a 45 minute period, to the latter

-54-

solution. The resulting reaction mixture was stored overnight in a cold room. Salts were filtered from the reaction mixture and the solution was evaporated. The residue was dissolved in EtOAc (1 L) and the insoluble  
5 solids were removed by filtration. The filtrate was washed with 1x300 mL of NaHCO<sub>3</sub> and 2x300 mL of saturated NaCl, dried over sodium sulfate and evaporated. The residue was triturated with EtOAc to give the title compound.

10        **2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine**

A solution of 3'-O-acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyl-4-triazoleuridine (103 g, 0.141 M) in dioxane (500 mL) and NH<sub>4</sub>OH (30 mL) was stirred at room temperature for 2 hours. The dioxane solution was  
15 evaporated and the residue azeotroped with MeOH (2x200 mL). The residue was dissolved in MeOH (300 mL) and transferred to a 2 liter stainless steel pressure vessel. MeOH (400 mL) saturated with NH<sub>3</sub> gas was added and the vessel heated to 100°C for 2 hours (TLC showed complete conversion). The  
20 vessel contents were evaporated to dryness and the residue was dissolved in EtOAc (500 mL) and washed once with saturated NaCl (200 mL). The organics were dried over sodium sulfate and the solvent was evaporated to give 85 g  
~~of the title compound.~~

25        **N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine**

2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine (85 g, 0.134 M) was dissolved in DMF (800 mL) and benzoic anhydride (37.2 g, 0.165 M) was added with  
30 stirring. After stirring for 3 hours, TLC showed the reaction to be approximately 95% complete. The solvent was evaporated and the residue azeotroped with MeOH (200 mL). The residue was dissolved in CHCl<sub>3</sub> (700 mL) and extracted with saturated NaHCO<sub>3</sub> (2x300 mL) and saturated NaCl (2x300  
35 mL), dried over MgSO<sub>4</sub> and evaporated to give a residue (96



-55-

g). The residue was chromatographed on a 1.5 kg silica column using EtOAc/hexane (1:1) containing 0.5% Et<sub>3</sub>NH as the eluting solvent. The pure product fractions were evaporated to give 90 g (90%) of the title compound.

5        **N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine-3'-amidite**

N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine (74 g, 0.10 M) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (1 L). Tetrazole diisopropylamine (7.1 g) and 2-cyanoethoxy-tetra-  
10 (isopropyl)phosphite (40.5 mL, 0.123 M) were added with stirring, under a nitrogen atmosphere. The resulting mixture was stirred for 20 hours at room temperature (TLC showed the reaction to be 95% complete). The reaction mixture was extracted with saturated NaHCO<sub>3</sub> (1x300 mL) and  
15 saturated NaCl (3x300 mL). The aqueous washes were back-extracted with CH<sub>2</sub>Cl<sub>2</sub> (300 mL), and the extracts were combined, dried over MgSO<sub>4</sub> and concentrated. The residue obtained was chromatographed on a 1.5 kg silica column using EtOAc/hexane (3:1) as the eluting solvent. The pure  
20 fractions were combined to give 90.6 g (87%) of the title compound.

**2'-O-(Aminoxyethyl) nucleoside amidites and 2'-O-(dimethylaminoxyethyl) nucleoside amidites**

**2'-(Dimethylaminoxyethoxy) nucleoside amidites**

25        2'-(Dimethylaminoxyethoxy) nucleoside amidites [also known in the art as 2'-O-(dimethylaminoxyethyl) nucleoside amidites] are prepared as described in the following paragraphs. Adenosine, cytidine and guanosine nucleoside amidites are prepared similarly to the thymidine (5-methyluridine) except the exocyclic amines are protected  
30 with a benzoyl moiety in the case of adenosine and cytidine and with isobutyryl in the case of guanosine.

-56-

5'-O-tert-Butyldiphenylsilyl-O<sup>2</sup>-2'-anhydro-5-methyluridine

O<sup>2</sup>-2'-anhydro-5-methyluridine (Pro. Bio. Sint., Varese, Italy, 100.0g, 0.416 mmol), dimethylaminopyridine (0.66g, 0.013eq, 0.0054mmol) were dissolved in dry pyridine (500 ml) at ambient temperature under an argon atmosphere and with mechanical stirring. tert-Butyldiphenylchlorosilane (125.8g, 119.0mL, 1.1eq, 0.458mmol) was added in one portion. The reaction was stirred for 16 h at ambient temperature. TLC (R<sub>f</sub> 0.22, ethyl acetate) indicated a complete reaction. The solution was concentrated under reduced pressure to a thick oil. This was partitioned between dichloromethane (1 L) and saturated sodium bicarbonate (2x1 L) and brine (1 L). The organic layer was dried over sodium sulfate and concentrated under reduced pressure to a thick oil. The oil was dissolved in a 1:1 mixture of ethyl acetate and ethyl ether (600mL) and the solution was cooled to -10°C. The resulting crystalline product was collected by filtration, washed with ethyl ether (3x200 mL) and dried (40°C, 1mm Hg, 24 h) to 149g (74.8%) of white solid. TLC and NMR were consistent with pure product.

~~5'-O-tert-Butyldiphenylsilyl-2'-O-(2-hydroxyethyl)-5-methyluridine~~

In a 2 L stainless steel, unstirred pressure reactor was added borane in tetrahydrofuran (1.0 M, 2.0 eq, 622 mL). In the fume hood and with manual stirring, ethylene glycol (350 mL, excess) was added cautiously at first until the evolution of hydrogen gas subsided. 5'-O-tert-Butyldiphenylsilyl-O<sup>2</sup>-2'-anhydro-5-methyluridine (149 g, 0.311 mol) and sodium bicarbonate (0.074 g, 0.003 eq) were added with manual stirring. The reactor was sealed and heated in an oil bath until an internal temperature of 160 °C was reached and then maintained for 16 h (pressure < 100

-57-

psig). The reaction vessel was cooled to ambient and opened. TLC (Rf 0.67 for desired product and Rf 0.82 for ara-T side product, ethyl acetate) indicated about 70% conversion to the product. In order to avoid additional side product formation, the reaction was stopped, concentrated under reduced pressure (10 to 1mm Hg) in a warm water bath (40-100°C) with the more extreme conditions used to remove the ethylene glycol. [Alternatively, once the low boiling solvent is gone, the remaining solution can be partitioned between ethyl acetate and water. The product will be in the organic phase.] The residue was purified by column chromatography (2kg silica gel, ethyl acetate-hexanes gradient 1:1 to 4:1). The appropriate fractions were combined, stripped and dried to product as a white crisp foam (84g, 50%), contaminated starting material (17.4g) and pure reusable starting material 20g. The yield based on starting material less pure recovered starting material was 58%. TLC and NMR were consistent with 99% pure product.

20        2'-O-([2-phthalimidoxy)ethyl]-5'-t-butylidiphenylsilyl-5-methyluridine

5'-O-tert-Butylidiphenylsilyl-2'-O-(2-hydroxyethyl)-5-methyluridine (20g, 36.98mmol) was mixed with triphenylphosphine (11.63g, 44.36mmol) and N-hydroxyphthalimide (7.24g, 44.36mmol). It was then dried over P<sub>2</sub>O<sub>5</sub> under high vacuum for two days at 40°C. The reaction mixture was flushed with argon and dry THF (369.8mL, Aldrich, sure seal bottle) was added to get a clear solution. Diethyl-azodicarboxylate (6.98mL, 44.36mmol) was added dropwise to the reaction mixture. The rate of addition is maintained such that resulting deep red coloration is just discharged before adding the next drop. After the addition was complete, the reaction was stirred for 4 hrs. By that time TLC showed the completion of the reaction (ethylacetate:hexane, 60:40). The solvent was

-58-

evaporated in vacuum. Residue obtained was placed on a flash column and eluted with ethyl acetate:hexane (60:40), to get 2'-O-([2-phthalimidooxy)ethyl]-5'-t-butyl-  
5 86%).

5'-O-tert-butyl-5-methyluridine-2'-O-[(2-formadoximinooxy)ethyl]-5-methyluridine

2'-O-([2-phthalimidooxy)ethyl]-5'-t-butyl-5-methyluridine (3.1g, 4.5mmol) was dissolved in dry CH<sub>2</sub>Cl<sub>2</sub>,  
10 (4.5mL) and methylhydrazine (300mL, 4.64mmol) was added dropwise at -10°C to 0°C. After 1 h the mixture was filtered, the filtrate was washed with ice cold CH<sub>2</sub>Cl<sub>2</sub> and the combined organic phase was washed with water, brine and dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>. The solution was concentrated  
15 to get 2'-O-(aminooxyethyl) thymidine, which was then dissolved in MeOH (67.5mL). To this formaldehyde (20% aqueous solution, w/w, 1.1 eq.) was added and the resulting mixture was stirred for 1 h. Solvent was removed under vacuum; residue chromatographed to get 5'-O-tert-  
20 butyl-5-methyluridine-2'-O-[(2-formadoximinooxy) ethyl]-5-methyluridine as white foam (1.95 g, 78%).

5'-O-tert-Butyl-5-methyluridine-2'-O-[N,N-

dimethylaminooxyethyl]-5-methyluridine

5'-O-tert-butyl-5-methyluridine-2'-O-[(2-formadoximinooxy)ethyl]-5-methyluridine (1.77g, 3.12mmol)  
25 was dissolved in a solution of 1M pyridinium p-toluenesulfonate (PPTS) in dry MeOH (30.6mL). Sodium cyanoborohydride (0.39g, 6.13mmol) was added to this solution at 10°C under inert atmosphere. The reaction  
30 mixture was stirred for 10 minutes at 10°C. After that the reaction vessel was removed from the ice bath and stirred at room temperature for 2 h, the reaction monitored by TLC (5% MeOH in CH<sub>2</sub>Cl<sub>2</sub>). Aqueous NaHCO<sub>3</sub> solution (5%, 10mL) was added and extracted with ethyl acetate (2x20mL). Ethyl

-59-

acetate phase was dried over anhydrous  $\text{Na}_2\text{SO}_4$ , evaporated to dryness. Residue was dissolved in a solution of 1M PPTS in MeOH (30.6mL). Formaldehyde (20% w/w, 30mL, 3.37mmol) was added and the reaction mixture was stirred at room temperature for 10 minutes. Reaction mixture cooled to 10°C in an ice bath, sodium cyanoborohydride (0.39g, 6.13mmol) was added and reaction mixture stirred at 10°C for 10 minutes. After 10 minutes, the reaction mixture was removed from the ice bath and stirred at room temperature for 2 hrs. To the reaction mixture 5%  $\text{NaHCO}_3$  (25mL) solution was added and extracted with ethyl acetate (2x25mL). Ethyl acetate layer was dried over anhydrous  $\text{Na}_2\text{SO}_4$  and evaporated to dryness. The residue obtained was purified by flash column chromatography and eluted with 5% MeOH in  $\text{CH}_2\text{Cl}_2$  to get 5'-O-tert-butylidiphenylsilyl-2'-O-[N,N-dimethylaminooxyethyl]-5-methyluridine as a white foam (14.6g, 80%).

**2'-O-(dimethylaminoxyethyl)-5-methyluridine**

Triethylamine trihydrofluoride (3.91mL, 24.0mmol) was dissolved in dry THF and triethylamine (1.67mL, 12mmol, dry, kept over KOH). This mixture of triethylamine-2HF was then added to 5'-O-tert-butylidiphenylsilyl-2'-O-[N,N-dimethylaminoxyethyl]-5-methyluridine (1.40g, 2.4mmol) and stirred at room temperature for 24 hrs. Reaction was monitored by TLC (5% MeOH in  $\text{CH}_2\text{Cl}_2$ ). Solvent was removed under vacuum and the residue placed on a flash column and eluted with 10% MeOH in  $\text{CH}_2\text{Cl}_2$  to get 2'-O-(dimethylaminoxyethyl)-5-methyluridine (766mg, 92.5%).

**5'-O-DMT-2'-O-(dimethylaminoxyethyl)-5-methyluridine**

2'-O-(dimethylaminoxyethyl)-5-methyluridine (750mg, 2.17mmol) was dried over  $\text{P}_2\text{O}_5$  under high vacuum overnight at 40°C. It was then co-evaporated with anhydrous pyridine (20mL). The residue obtained was dissolved in pyridine (11mL) under argon atmosphere. 4-dimethylaminopyridine (26.5mg, 2.60mmol), 4,4'-dimethoxytrityl chloride (880mg,

-60-

2.60mmol) was added to the mixture and the reaction mixture was stirred at room temperature until all of the starting material disappeared. Pyridine was removed under vacuum and the residue chromatographed and eluted with 10% MeOH in 5 CH<sub>2</sub>Cl<sub>2</sub> (containing a few drops of pyridine) to get 5'-O-DMT-2'-O-(dimethylamino-oxyethyl)-5-methyluridine (1.13g, 80%).

5'-O-DMT-2'-O-(2-N,N-dimethylaminooxyethyl)-5-methyluridine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite]

10 5'-O-DMT-2'-O-(dimethylaminooxyethyl)-5-methyluridine (1.08g, 1.67mmol) was co-evaporated with toluene (20mL). To the residue N,N-diisopropylamine tetrazonide (0.29g, 1.67mmol) was added and dried over P<sub>2</sub>O<sub>5</sub> under high vacuum overnight at 40°C. Then the reaction mixture was dissolved 15 in anhydrous acetonitrile (8.4mL) and 2-cyanoethyl-N,N,N<sup>1</sup>,N<sup>1</sup>-tetraisopropylphosphoramidite (2.12mL, 6.08mmol) was added. The reaction mixture was stirred at ambient temperature for 4 hrs under inert atmosphere. The progress of the reaction was monitored by TLC (hexane:ethyl acetate 20 1:1). The solvent was evaporated, then the residue was dissolved in ethyl acetate (70mL) and washed with 5% aqueous NaHCO<sub>3</sub> (40mL). Ethyl acetate layer was dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and concentrated. Residue obtained was chromatographed (ethyl acetate as eluent) to get 5'-O-DMT-25 2'-O-(2-N,N-dimethylaminooxyethyl)-5-methyluridine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite] as a foam (1.04g, 74.9%).

2'-(Aminooxyethoxy) nucleoside amidites

2'-(Aminooxyethoxy) nucleoside amidites [also known in 30 the art as 2'-O-(aminooxyethyl) nucleoside amidites] are prepared as described in the following paragraphs. Adenosine, cytidine and thymidine nucleoside amidites are prepared similarly.

-61-

N2-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite]

- The 2'-O-aminooxyethyl guanosine analog may be
- 5 obtained by selective 2'-O-alkylation of diaminopurine riboside. Multigram quantities of diaminopurine riboside may be purchased from Schering AG (Berlin) to provide 2'-O-(2-ethylacetyl) diaminopurine riboside along with a minor amount of the 3'-O-isomer. 2'-O-(2-ethylacetyl)
- 10 diaminopurine riboside may be resolved and converted to 2'-O-(2-ethylacetyl)guanosine by treatment with adenosine deaminase. (McGee, D. P. C., Cook, P. D., Guinosso, C. J., WO 94/02501 A1 940203.) Standard protection procedures should afford 2'-O-(2-ethylacetyl)-5'-O-(4,4'-
- 15 dimethoxytrityl)guanosine and 2-N-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine which may be reduced to provide 2-N-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine. As before the
- 20 hydroxyl group may be displaced by N-hydroxyphthalimide via a Mitsunobu reaction, and the protected nucleoside may phosphitylated as usual to yield 2-N-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine-3'-[(2-cyanoethyl)-N,N-
- 25 diisopropylphosphoramidite].

**2'-dimethylaminoethoxyethoxy (2'-DMAEOE) nucleoside amidites**

- 2'-dimethylaminoethoxyethoxy nucleoside amidites (also known in the art as 2'-O-dimethylaminoethoxyethyl, i.e.,
- 30 2'-O-CH<sub>2</sub>-O-CH<sub>2</sub>-N(CH<sub>3</sub>)<sub>2</sub>, or 2'-DMAEOE nucleoside amidites) are prepared as follows. Other nucleoside amidites are prepared similarly.

-62-

**2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyl uridine**

2[2-(Dimethylamino)ethoxy]ethanol (Aldrich, 6.66 g, 50 mmol) is slowly added to a solution of borane in tetrahydrofuran (1 M, 10 mL, 10 mmol) with stirring in a 100 mL bomb. Hydrogen gas evolves as the solid dissolves. O<sup>2</sup>-,2'-anhydro-5-methyluridine (1.2 g, 5 mmol), and sodium bicarbonate (2.5 mg) are added and the bomb is sealed, placed in an oil bath and heated to 155°C for 26 hours. The bomb is cooled to room temperature and opened. The crude solution is concentrated and the residue partitioned between water (200 mL) and hexanes (200 mL). The excess phenol is extracted into the hexane layer. The aqueous layer is extracted with ethyl acetate (3x200 mL) and the combined organic layers are washed once with water, dried over anhydrous sodium sulfate and concentrated. The residue is columned on silica gel using methanol/methylene chloride 1:20 (which has 2% triethylamine) as the eluent. As the column fractions are concentrated a colorless solid forms which is collected to give the title compound as a white solid.

**5'-O-dimethoxytrityl-2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyl uridine**

To 0.5 g (1.3 mmol) of 2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyl uridine in anhydrous pyridine (8 mL), triethylamine (0.36 mL) and dimethoxytrityl chloride (DMT-Cl, 0.87 g, 2 eq.) are added and stirred for 1 hour. The reaction mixture is poured into water (200 mL) and extracted with CH<sub>2</sub>Cl<sub>2</sub> (2x200 mL). The combined CH<sub>2</sub>Cl<sub>2</sub> layers are washed with saturated NaHCO<sub>3</sub> solution, followed by saturated NaCl solution and dried over anhydrous sodium sulfate. Evaporation of the solvent followed by silica gel chromatography using MeOH:CH<sub>2</sub>Cl<sub>2</sub>:Et<sub>3</sub>N (20:1, v/v, with 1% triethylamine) gives the title compound.



-63-

5'-O-Dimethoxytrityl-2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl)]-5-methyl uridine-3'-O-(cyanoethyl-N,N-diisopropyl)phosphoramidite

Diisopropylaminotetrazolide (0.6 g) and 2-cyanoethoxy-  
5 N,N-diisopropyl phosphoramidite (1.1 mL, 2 eq.) are added to a solution of 5'-O-dimethoxytrityl-2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl)]-5-methyluridine (2.17 g, 3 mmol) dissolved in CH<sub>2</sub>Cl<sub>2</sub> (20 mL) under an atmosphere of argon. The reaction mixture is stirred overnight and the  
10 solvent evaporated. The resulting residue is purified by silica gel flash column chromatography with ethyl acetate as the eluent to give the title compound.

#### Example 2

##### Oligonucleotide synthesis

15 Unsubstituted and substituted phosphodiester (P=O) oligonucleotides are synthesized on an automated DNA synthesizer (Applied Biosystems model 380B) using standard phosphoramidite chemistry with oxidation by iodine.

Phosphorothioates (P=S) are synthesized as for the  
20 phosphodiester oligonucleotides except the standard oxidation bottle was replaced by 0.2 M solution of 3H-1,2-benzodithiole-3-one 1,1-dioxide in acetonitrile for the stepwise thiation of the phosphite linkages. The thiation wait step was increased to 68 sec and was followed by the  
25 capping step. After cleavage from the CPG column and deblocking in concentrated ammonium hydroxide at 55°C (18 h), the oligonucleotides were purified by precipitating twice with 2.5 volumes of ethanol from a 0.5 M NaCl solution. Phosphinate oligonucleotides are prepared as  
30 described in U.S. Patent 5,508,270, herein incorporated by reference.

Alkyl phosphonate oligonucleotides are prepared as described in U.S. Patent 4,469,863, herein incorporated by reference.

-64-

3'-Deoxy-3'-methylene phosphonate oligonucleotides are prepared as described in U.S. Patents 5,610,289 or 5,625,050, herein incorporated by reference.

Phosphoramidite oligonucleotides are prepared as described in U.S. Patent, 5,256,775 or U.S. Patent 5,366,878, herein incorporated by reference.

Alkylphosphonothioate oligonucleotides are prepared as described in published PCT applications PCT/US94/00902 and PCT/US93/06976 (published as WO 94/17093 and WO 94/02499, respectively), herein incorporated by reference.

3'-Deoxy-3'-amino phosphoramidate oligonucleotides are prepared as described in U.S. Patent 5,476,925, herein incorporated by reference.

Phosphotriester oligonucleotides are prepared as described in U.S. Patent 5,023,243, herein incorporated by reference.

Borano phosphate oligonucleotides are prepared as described in U.S. Patents 5,130,302 and 5,177,198, both herein incorporated by reference.

### Example 3

#### Oligonucleoside Synthesis

Methylenemethylimino linked oligonucleosides, also identified as MMI linked oligonucleosides, methylenedimethylhydrazo linked oligonucleosides, also identified as MDH linked oligonucleosides, and methylenecarbonylamino linked oligonucleosides, also identified as amide-3 linked oligonucleosides, and methyleneaminocarbonyl linked oligonucleosides, also identified as amide-4 linked oligonucleosides, as well as mixed backbone compounds having, for instance, alternating MMI and P=O or P=S linkages are prepared as described in U.S. Patents 5,378,825, 5,386,023, 5,489,677, 5,602,240 and 5,610,289, all of which are herein incorporated by reference.

Formacetal and thioformacetal linked oligonucleosides are prepared as described in U.S. Patents 5,264,562 and 5,264,564, herein incorporated by reference.

Ethylene oxide linked oligonucleosides are prepared as described in U.S. Patent 5,223,618, herein incorporated by reference.

#### Example 4

##### PNA Synthesis

Peptide nucleic acids (PNAs) are prepared in accordance with any of the various procedures referred to in *Peptide Nucleic Acids (PNA): Synthesis, Properties and Potential Applications, Bioorganic & Medicinal Chemistry*, 1996, 4, 5-23. They may also be prepared in accordance with U.S. Patents 5,539,082, 5,700,922, and 5,719,262, herein incorporated by reference.

#### Example 5

##### Synthesis of Chimeric Oligonucleotides

Chimeric oligonucleotides, oligonucleosides or mixed oligonucleotides/oligonucleosides of the invention can be of several different types. These include a first type wherein the "gap" segment of linked nucleosides is positioned between 5' and 3' "wing" segments of linked nucleosides and a second "open end" type wherein the "gap" segment is located at either the 3' or the 5' terminus of the oligomeric compound. Oligonucleotides of the first type are also known in the art as "gapmers" or gapped oligonucleotides. Oligonucleotides of the second type are also known in the art as "hemimers" or "wingmers".

##### [2'-O-Me]--[2'-deoxy]--[2'-O-Me] Chimeric

#### 30 Phosphorothioate Oligonucleotides

Chimeric oligonucleotides having 2'-O-alkyl phosphorothioate and 2'-deoxy phosphorothioate oligonucleotide segments are synthesized using an Applied Biosystems automated DNA synthesizer Model 380B, as above.

Oligonucleotides are synthesized using the automated synthesizer and 2'-deoxy-5'-dimethoxytrityl-3'-O-phosphoramidite for the DNA portion and 5'-dimethoxytrityl-2'-O-methyl-3'-O-phosphoramidite for 5' and 3' wings. The standard synthesis cycle is modified by increasing the wait step after the delivery of tetrazole and base to 600 s repeated four times for RNA and twice for 2'-O-methyl. The fully protected oligonucleotide is cleaved from the support and the phosphate group is deprotected in 3:1 ammonia/ethanol at room temperature overnight then lyophilized to dryness. Treatment in methanolic ammonia for 24 hrs at room temperature is then done to deprotect all bases and sample was again lyophilized to dryness. The pellet is resuspended in 1M TBAF in THF for 24 hrs at room temperature to deprotect the 2' positions. The reaction is then quenched with 1M TEAA and the sample is then reduced to 1/2 volume by rotovac before being desalted on a G25 size exclusion column. The oligo recovered is then analyzed spectrophotometrically for yield and for purity by capillary electrophoresis and by mass spectrometry.

[2'-O-(2-Methoxyethyl)]--[2'-deoxy]--[2'-O-(Methoxyethyl)] Chimeric Phosphorothioate Oligonucleotides

[2'-O-(2-methoxyethyl)]--[2'-deoxy]--[2'-O-(methoxyethyl)] chimeric phosphorothioate oligonucleotides were prepared as per the procedure above for the 2'-O-methyl chimeric oligonucleotide, with the substitution of 2'-O-(methoxyethyl) amidites for the 2'-O-methyl amidites.

[2'-O-(2-Methoxyethyl)Phosphodiester]--[2'-deoxy Phosphorothioate]--[2'-O-(2-Methoxyethyl) Phosphodiester] Chimeric Oligonucleotides

[2'-O-(2-methoxyethyl phosphodiester)]--[2'-deoxy phosphorothioate]--[2'-O-(methoxyethyl) phosphodiester] chimeric oligonucleotides are prepared as per the above

-67-

procedure for the 2'-O-methyl chimeric oligonucleotide with the substitution of 2'-O-(methoxyethyl) amidites for the 2'-O-methyl amidites, oxidization with iodine to generate the phosphodiester internucleotide linkages within the wing portions of the chimeric structures and sulfurization utilizing 3,4-dihydro-2H-benzothiole-3-one 1,1-dioxide (Beaucage Reagent) to generate the phosphorothioate internucleotide linkages for the center gap.

Other chimeric oligonucleotides, chimeric oligonucleosides and mixed chimeric oligonucleotides/oligonucleosides are synthesized according to United States patent 5,623,065, herein incorporated by reference.

#### Example 6

##### Oligonucleotide Isolation

After cleavage from the controlled pore glass column (Applied Biosystems) and deblocking in concentrated ammonium hydroxide at 55°C for 18 hours, the oligonucleotides or oligonucleosides are purified by precipitation twice out of 0.5 M NaCl with 2.5 volumes ethanol. Synthesized oligonucleotides were analyzed by polyacrylamide gel electrophoresis on denaturing gels and judged to be at least 85% full length material. The relative amounts of phosphorothioate and phosphodiester linkages obtained in synthesis were periodically checked by <sup>31</sup>P nuclear magnetic resonance spectroscopy, and for some studies oligonucleotides were purified by HPLC, as described by Chiang et al., *J. Biol. Chem.* 1991, 266, 18162-18171. Results obtained with HPLC-purified material were similar to those obtained with non-HPLC purified material.

#### Example 7

##### Oligonucleotide Synthesis - 96 Well Plate Format

Oligonucleotides were synthesized via solid phase P(III) phosphoramidite chemistry on an automated

-68-

synthesizer capable of assembling 96 sequences simultaneously in a standard 96 well format. Phosphodiester internucleotide linkages were afforded by oxidation with aqueous iodine. Phosphorothioate internucleotide linkages were generated by sulfurization utilizing 3,4-dihydro-2H-benzothiole-3-one 1,1-dioxide (Beaucage Reagent) in anhydrous acetonitrile. Standard base-protected beta-cyanoethyl-diisopropyl phosphoramidites were purchased from commercial vendors (e.g. PE-Applied Biosystems, Foster City, CA, or Pharmacia, Piscataway, NJ). Non-standard nucleosides are synthesized as per known literature or patented methods. They are utilized as base-protected beta-cyanoethyl-diisopropyl phosphoramidites.

Oligonucleotides were cleaved from support and deprotected with concentrated  $\text{NH}_4\text{OH}$  at elevated temperature ( $55-60^\circ\text{C}$ ) for 12-16 hours and the released product then dried in vacuo. The dried product was then re-suspended in sterile water to afford a master plate from which all analytical and test plate samples are then diluted utilizing robotic pipettors.

#### Example 8

##### Oligonucleotide Analysis - 96 Well Plate Format

The concentration of oligonucleotide in each well was ~~measured by dilution of samples and UV absorption~~ spectroscopy. The full-length integrity of the individual products was evaluated by capillary electrophoresis (CE) in either the 96 well format (Beckman P/ACE™ MDQ) or, for individually prepared samples, on a commercial CE apparatus (e.g., Beckman P/ACE™ 5000, ABI 270). Base and backbone composition was confirmed by mass analysis of the compounds utilizing electrospray-mass spectroscopy. All assay test plates were diluted from the master plate using single and multi-channel robotic pipettors. Plates were judged to be acceptable if at least 85% of the compounds on the plate were at least 85% full length.

**Example 9****Cell culture and oligonucleotide treatment**

The effect of antisense compounds on target nucleic acid expression can be tested in any of a variety of cell types provided that the target nucleic acid is present at measurable levels. This can be routinely determined using, for example, PCR or Northern blot analysis. The following four cell types are provided for illustrative purposes, but other cell types can be routinely used.

10

**T-24 cells:**

The transitional cell bladder carcinoma cell line T-24 was obtained from the American Type Culture Collection (ATCC) (Manassas, VA). T-24 cells were routinely cultured in complete McCoy's 5A basal media (Gibco/Life Technologies, Gaithersburg, MD) supplemented with 10% fetal calf serum (Gibco/Life Technologies, Gaithersburg, MD), penicillin 100 units per mL, and streptomycin 100 micrograms per mL (Gibco/Life Technologies, Gaithersburg, MD). Cells were routinely passaged by trypsinization and dilution when they reached 90% confluence. Cells were seeded into 96-well plates (Falcon-Primaria #3872) at a density of 7000 cells/well for use in RT-PCR analysis.

For Northern blotting or other analysis, cells may be seeded onto 100 mm or other standard tissue culture plates and treated similarly, using appropriate volumes of medium and oligonucleotide.

**A549 cells:**

The human lung carcinoma cell line A549 was obtained from the American Type Culture Collection (ATCC) (Manassas, VA). A549 cells were routinely cultured in DMEM basal media (Gibco/Life Technologies, Gaithersburg, MD) supplemented with 10% fetal calf serum (Gibco/Life Technologies, Gaithersburg, MD), penicillin 100 units per

-70-

mL, and streptomycin 100 micrograms per mL (Gibco/Life Technologies, Gaithersburg, MD). Cells were routinely passaged by trypsinization and dilution when they reached 90% confluence.

5

#### NHDF cells:

Human neonatal dermal fibroblast (NHDF) were obtained from the Clonetics Corporation (Walkersville MD). NHDFs were routinely maintained in Fibroblast Growth Medium (Clonetics Corporation, Walkersville MD) supplemented as recommended by the supplier. Cells were maintained for up to 10 passages as recommended by the supplier.

#### HEK cells:

15 Human embryonic keratinocytes (HEK) were obtained from the Clonetics Corporation (Walkersville MD). HEKs were routinely maintained in Keratinocyte Growth Medium (Clonetics Corporation, Walkersville MD) formulated as recommended by the supplier. Cells were routinely  
20 maintained for up to 10 passages as recommended by the supplier.

#### Treatment with antisense compounds:

---

when cells reached 80% confluency, they were treated  
25 with oligonucleotide. For cells grown in 96-well plates, wells were washed once with 200  $\mu$ L OPTI-MEM™-1 reduced-serum medium (Gibco BRL) and then treated with 130  $\mu$ L of OPTI-MEM™-1 containing 3.75  $\mu$ g/mL LIPOFECTIN™ (Gibco BRL) and the desired oligonucleotide at a final concentration of  
30 150 nM. After 4 hours of treatment, the medium was replaced with fresh medium. Cells were harvested 16 hours after oligonucleotide treatment.



-71-

**Example 10****Analysis of oligonucleotide inhibition of PI3 kinase p110 beta expression****Antisense modulation of PI3 kinase p110 beta**

5 expression can be assayed in a variety of ways known in the art. For example, PI3 kinase p110 beta mRNA levels can be quantitated by, e.g., Northern blot analysis, competitive polymerase chain reaction (PCR), or real-time PCR (RT-PCR). Real-time quantitative PCR is presently preferred. RNA  
10 analysis can be performed on total cellular RNA or poly(A)+ mRNA. Methods of RNA isolation are taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 1, pp. 4.1.1-4.2.9 and 4.5.1-4.5.3, John Wiley & Sons, Inc., 1993. Northern blot analysis is  
15 routine in the art and is taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 1, pp. 4.2.1-4.2.9, John Wiley & Sons, Inc., 1996. Real-time quantitative (PCR) can be conveniently accomplished using the commercially available ABI PRISM™ 7700 Sequence  
20 Detection System, available from PE-Applied Biosystems, Foster City, CA and used according to manufacturer's instructions. Other methods of PCR are also known in the art.

PI3 kinase p110 beta protein levels can be quantitated  
25 in a variety of ways well known in the art, such as immunoprecipitation, Western blot analysis (immunoblotting), ELISA or fluorescence-activated cell sorting (FACS). Antibodies directed to PI3 kinase p110 beta can be identified and obtained from a variety of  
30 sources, such as the MSRS catalog of antibodies (Aerie Corporation, Birmingham, MI), or can be prepared via conventional antibody generation methods. Methods for preparation of polyclonal antisera are taught in, for example, Ausubel, F.M. et al., *Current Protocols in*

-72-

- Molecular Biology, Volume 2, pp. 11.12.1-11.12.9, John Wiley & Sons, Inc., 1997. Preparation of monoclonal antibodies is taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 5 11.4.1-11.11.5, John Wiley & Sons, Inc., 1997.

- Immunoprecipitation methods are standard in the art and can be found at, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 10.16.1-10.16.11, John Wiley & Sons, Inc., 1998. Western blot (immunoblot) analysis is standard in the art and can be found at, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 10.8.1-10.8.21, John Wiley & Sons, Inc., 1997. Enzyme-linked immunosorbent assays (ELISA) are standard in the art and 15 can be found at, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 11.2.1-11.2.22, John Wiley & Sons, Inc., 1991.

#### Example 11

##### Poly(A)+ mRNA isolation

- 20 Poly(A)+ mRNA was isolated according to Miura et al., *Clin. Chem.*, 1996, 42, 1758-1764. Other methods for poly(A)+ mRNA isolation are taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 1, pp. 4.5.1-4.5.3, John Wiley & Sons, Inc., 1993. Briefly, for cells grown on 96-well plates, 25 growth medium was removed from the cells and each well was washed with 200  $\mu$ L cold PBS. 60  $\mu$ L lysis buffer (10 mM Tris-HCl, pH 7.6, 1 mM EDTA, 0.5 M NaCl, 0.5% NP-40, 20 mM vanadyl-ribonucleoside complex) was added to each well, the 30 plate was gently agitated and then incubated at room temperature for five minutes. 55  $\mu$ L of lysate was transferred to Oligo d(T) coated 96-well plates (AGCT Inc., Irvine CA). Plates were incubated for 60 minutes at room temperature, washed 3 times with 200  $\mu$ L of wash buffer (10

-73-

mM Tris-HCl pH 7.6, 1 mM EDTA, 0.3 M NaCl). After the final wash, the plate was blotted on paper towels to remove excess wash buffer and then air-dried for 5 minutes. 60  $\mu$ L of elution buffer (5 mM Tris-HCl pH 7.6), preheated to 70°C was added to each well, the plate was incubated on a 90°C hot plate for 5 minutes, and the eluate was then transferred to a fresh 96-well plate.

Cells grown on 100 mm or other standard plates may be treated similarly, using appropriate volumes of all solutions.

#### Example 12

##### Total RNA Isolation

Total mRNA was isolated using an RNEASY 96™ kit and buffers purchased from Qiagen Inc. (Valencia CA) following the manufacturer's recommended procedures. Briefly, for cells grown on 96-well plates, growth medium was removed from the cells and each well was washed with 200  $\mu$ L cold PBS. 100  $\mu$ L Buffer RLT was added to each well and the plate vigorously agitated for 20 seconds. 100  $\mu$ L of 70% ethanol was then added to each well and the contents mixed by pipetting three times up and down. The samples were then transferred to the RNEASY 96™ well plate attached to a QIAVAC™ manifold fitted with a waste collection tray and attached to a vacuum source. Vacuum was applied for 15 seconds. 1 mL of Buffer RW1 was added to each well of the RNEASY 96™ plate and the vacuum again applied for 15 seconds. 1 mL of Buffer RPE was then added to each well of the RNEASY 96™ plate and the vacuum applied for a period of 15 seconds. The Buffer RPE wash was then repeated and the vacuum was applied for an additional 10 minutes. The plate was then removed from the QIAVAC™ manifold and blotted dry on paper towels. The plate was then re-attached to the QIAVAC™ manifold fitted with a collection tube rack containing 1.2 mL collection tubes. RNA was then eluted by pipetting 60  $\mu$ L water into each well, incubating 1 minute,

-74-

and then applying the vacuum for 30 seconds. The elution step was repeated with an additional 60  $\mu$ L water.

The repetitive pipetting and elution steps may be automated using a QIAGEN Bio-Robot 9604 (Qiagen, Inc., Valencia CA). Essentially after lysing of the cells on the culture plate, the plate is transferred to the robot deck where the pipetting, DNase treatment and elution steps are carried out.

#### Example 13

#### 10 Real-time Quantitative PCR Analysis of PI3 kinase p110 beta mRNA Levels

Quantitation of PI3 kinase p110 beta mRNA levels was determined by real-time quantitative PCR using the ABI PRISM™ 7700 Sequence Detection System (PE-Applied Biosystems, Foster City, CA) according to manufacturer's instructions. This is a closed-tube, non-gel-based, fluorescence detection system which allows high-throughput quantitation of polymerase chain reaction (PCR) products in real-time. As opposed to standard PCR, in which amplification products are quantitated after the PCR is completed, products in real-time quantitative PCR are quantitated as they accumulate. This is accomplished by including in the PCR reaction an oligonucleotide probe that anneals specifically between the forward and reverse PCR primers, and contains two fluorescent dyes. A reporter dye (e.g., JOE or FAM, obtained from either Operon Technologies Inc., Alameda, CA or PE-Applied Biosystems, Foster City, CA) is attached to the 5' end of the probe and a quencher dye (e.g., TAMRA, obtained from either Operon Technologies Inc., Alameda, CA or PE-Applied Biosystems, Foster City, CA) is attached to the 3' end of the probe. When the probe and dyes are intact, reporter dye emission is quenched by the proximity of the 3' quencher dye. During amplification, annealing of the probe to the target sequence creates a substrate that can be cleaved by the 5'-

-75-

exonuclease activity of Taq polymerase. During the extension phase of the PCR amplification cycle, cleavage of the probe by Taq polymerase releases the reporter dye from the remainder of the probe (and hence from the quencher moiety) and a sequence-specific fluorescent signal is generated. With each cycle, additional reporter dye molecules are cleaved from their respective probes, and the fluorescence intensity is monitored at regular intervals by laser optics built into the ABI PRISM™ 7700 Sequence Detection System. In each assay, a series of parallel reactions containing serial dilutions of mRNA from untreated control samples generates a standard curve that is used to quantitate the percent inhibition after antisense oligonucleotide treatment of test samples.

PCR reagents were obtained from PE-Applied Biosystems, Foster City, CA. RT-PCR reactions were carried out by adding 25 µL PCR cocktail (1x TAQMAN™ buffer A, 5.5 mM MgCl<sub>2</sub>, 300 µM each of dATP, dCTP and dGTP, 600 µM of dUTP, 100 nM each of forward primer, reverse primer, and probe, 20 Units RNase inhibitor, 1.25 Units AMPLITAQ GOLD™, and 12.5 Units MuLV reverse transcriptase) to 96 well plates containing 25 µL poly(A) mRNA solution. The RT reaction was carried out by incubation for 30 minutes at 48°C. Following a 10 minute incubation at 95°C to activate the AMPLITAQ GOLD™, 40 cycles of a two-step PCR protocol were carried out: 95°C for 15 seconds (denaturation) followed by 60°C for 1.5 minutes (annealing/extension). PI3 kinase p110 beta probes and primers were designed to hybridize to the human PI3 kinase p110 beta sequence, using published sequence information (GenBank accession number S67334, incorporated herein as SEQ ID NO:1).

For PI3 kinase p110 beta the PCR primers were:  
forward primer: GCGGTGGATTACAGATAGCA (SEQ ID NO: 2)  
reverse primer: TGGTAGCTTCCCGAGGTACCT (SEQ ID NO: 3) and  
the PCR probe was: FAM-TGGATTTCTTTTGCCCACTGGGA-TAMRA

-76-

(SEQ ID NO: 4) where FAM (PE-Applied Biosystems, Foster City, CA) is the fluorescent reporter dye) and TAMRA (PE-Applied Biosystems, Foster City, CA) is the quencher dye.

For GAPDH the PCR primers were:

- 5 forward primer: GAAGGTGAAGGTCGGAGTC (SEQ ID NO: 5)  
reverse primer: GAAGATGGTGATGGGATTTC (SEQ ID NO: 6) and the  
PCR probe was: 5' JOE-CAAGCTTCCCGTTCTCAGCC- TAMRA 3' (SEQ  
ID NO: 7) where JOE (PE-Applied Biosystems, Foster City,  
CA) is the fluorescent reporter dye) and TAMRA (PE-Applied  
10 Biosystems, Foster City, CA) is the quencher dye.

#### Example 14

##### Northern blot analysis of PI3 kinase p110 beta mRNA levels

- Eighteen hours after antisense treatment, cell  
monolayers were washed twice with cold PBS and lysed in 1  
15 mL RNAzol™ (TEL-TEST "B" Inc., Friendswood, TX). Total RNA  
was prepared following manufacturer's recommended  
protocols. Twenty micrograms of total RNA was fractionated  
by electrophoresis through 1.2% agarose gels containing  
1.1% formaldehyde using a MOPS buffer system (AMRESCO, Inc.  
20 Solon, OH). RNA was transferred from the gel to HYBOND™-N+  
nylon membranes (Amersham Pharmacia Biotech, Piscataway,  
NJ) by overnight capillary transfer using a  
~~Northern/Southern Transfer buffer system (TEL-TEST "B"~~  
Inc., Friendswood, TX). RNA transfer was confirmed by UV  
25 visualization. Membranes were fixed by UV cross-linking  
using a STRATALINKER™ UV Crosslinker 2400 (Stratagene,  
Inc, La Jolla, CA).

- Membranes were probed using QUICKHYB™ hybridization  
solution (Stratagene, La Jolla, CA) using manufacturer's  
30 recommendations for stringent conditions with a PI3 kinase  
p110 beta specific probe prepared by PCR using the forward  
primer GCGGTGGATTACAGATAGCA (SEQ ID NO: 2) and the reverse  
primer TGGTAGCTTCCCGAGGTACCT (SEQ ID NO: 3). To normalize  
for variations in loading and transfer efficiency membranes  
35 were stripped and probed for glyceraldehyde-3-phosphate

-77-

dehydrogenase (GAPDH) RNA (Clontech, Palo Alto, CA). Hybridized membranes were visualized and quantitated using a PHOSPHORIMAGER™ and IMAGEQUANT™ Software V3.3 (Molecular Dynamics, Sunnyvale, CA). Data was normalized to GAPDH levels in untreated controls.

**Example 15****Antisense inhibition of PI3 kinase p110 beta expression-phosphorothioate oligodeoxynucleotides**

In accordance with the present invention, a series of oligonucleotides were designed to target different regions of the human PI3 kinase p110 beta RNA, using published sequences (GenBank accession number S67334, incorporated herein as SEQ ID NO: 1). The oligonucleotides are shown in Table 1. Target sites are indicated by nucleotide numbers, as given in the sequence source reference (Genbank accession no. S67334), to which the oligonucleotide binds. All compounds in Table 1 are oligodeoxynucleotides with phosphorothioate backbones (internucleoside linkages) throughout. The compounds were analyzed for effect on PI3 kinase p110 beta mRNA levels by quantitative real-time PCR as described in other examples herein. Data are averages from two experiments. If present, "N.D." indicates "no data".

Table 1

Inhibition of PI3 kinase p110 beta mRNA levels by phosphorothioate oligodeoxynucleotides

ISIS#	REGION	TARGET SITE	SEQUENCE	% Inhib.	SEQ ID NO.
31960	Start Codon	2	cattatgaaactgaagcaca	0	8
30 31961	Coding	4	ggcattatgaaactgaagca	0	9
31962	Coding	23	gatgtctgccatagcaggag	0	10
31963	Coding	41	caccgcccagatgtcaagga	50	11
31964	Coding	56	tgctatctgtgaatccaccg	1	12
31965	Coding	74	aggtatggagccatcagatg	0	13
35 31966	Coding	92	gggcaaaaggaaatccacag	0	14
31967	Coding	111	actggatataaatcccagtg	0	15

-78-

	31968	Coding	128	ttccccgaggtacctccaact	30	16
	31969	Coding	144	tataagaaatggtagcttcc	0	17
	31970	Coding	161	ccataacatctgcttaatat	0	18
	31971	Coding	177	aattgtgaacttgcttccat	26	19
5	31972	Coding	194	gaggttgaacattgggtaat	21	20
	31973	Coding	209	gtcaatatccataaggaggt	0	21
	31974	Coding	225	atgcaaacatataggagtca	0	22
	31975	Coding	303	ctggaagaaaaggtctgaca	0	23
	31976	Coding	408	gaccccttcaaggaatcaa	9	24
10	31977	Coding	544	aagttttcagggatggatgg	0	25
	31978	Coding	587	atgaacagctacgatgagct	27	26
	31979	Coding	612	taaacacgtcctggcagttt	0	27
	31980	Coding	635	catattaggagacacvtgaa	0	28
	31981	Coding	662	tgccaattcatttactttga	0	29
15	31982	Coding	692	cttcccatgaatagtcaa	47	30
	31983	Coding	727	tgcaacacataatcataggg	0	31
	31984	Coding	772	attagtggatgatcacaaa	1	32
	31985	Coding	800	cacacagttccggatatact	0	33
	31986	Coding	859	tacattttcttgatcttgca	0	34
20	31987	Coding	1049	accagccctgacatgaactt	55	35
	31988	Coding	1360	tgtaatattatgtctccagt	10	36
	31989	Coding	1777	ggccaaatctgaagcagcgc	0	37
	31990	Coding	1894	tattgagaaagttcttcac	14	38
	31991	Coding	2132	attgagtgttcaacctgct	0	39
25	31992	Coding	2285	gaggataacacatgggttca	0	40
	31993	Coding	2440	aacatttggagtgtcaacat	0	41
	31994	Coding	2497	taaggcaacatccgaagatc	4	42
	31995	Coding	2706	agtagccagcacaggacagt	0	43
	31996	Coding	2747	actatgtctgtcaccaatcc	48	44
30	31997	Coding	2975	attcccatgccgtcgtaaaa	0	45
	31998	Coding	2999	cagcccaaagagagtga	0	46
	31999	Coding	3154	ttcacttttagtagtccagt	11	47

As shown in Table 1, SEQ ID NOs 11, 16, 19, 20, 26, 30, 35 and 44 demonstrated at least 20% inhibition of PI3 kinase p110 beta expression in this assay and are therefore preferred.

#### Example 16:

Antisense inhibition of PI3 kinase p110 beta expression-  
phosphorothioate 2'-MOE gapmer oligonucleotides

In accordance with the present invention, a second series of oligonucleotides targeted to human PI3 kinase p110 beta were synthesized. The oligonucleotide sequences



-79-

are shown in Table 2. Target sites are indicated by nucleotide numbers, as given in the sequence source reference (Genbank accession no. S67334), to which the oligonucleotide binds.

- 5 All compounds in Table 2 are chimeric oligonucleotides ("gapmers") 20 nucleotides in length, composed of a central "gap" region consisting of ten 2'-deoxynucleotides, which is flanked on both sides (5' and 3' directions) by five-nucleotide "wings". The wings are composed of 2'-
- 10 methoxyethyl (2'-MOE)nucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotide. Cytidine residues in the 2'-MOE wings are 5-methylcytidines.

Data were obtained by real-time quantitative PCR as

15 described in other examples herein and are averaged from two experiments. If present, "N.D." indicates "no data".

Table 2

Inhibition of PI3 kinase p110 beta mRNA levels by chimeric

20 phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap

	ISIS#	REGION	TARGET	SEQUENCE	% Inhib.	SEQ ID NO.
			SITE			
	32000	Start Codon	2	cattatgaaactgaagcaca	77	8
25	32001	Coding	4	ggcattatgaaactgaagca	24	9
	32002	Coding	23	gatgtctgccatagcaggag	54	10
	32003	Coding	41	caccgcccagatgtcaagga	82	11
	32004	Coding	56	tgctatctgtgaatccaccg	87	12
	32005	Coding	74	aggtatggagccatcagatg	80	13
30	32006	Coding	92	gggcaaaaggaaatccacag	70	14
	32007	Coding	111	actggatataaatcccagtg	0	15
	32008	Coding	128	ttcccaggtacctccaact	85	16
	32009	Coding	144	tataagaaatggtagcttcc	37	17
	32010	Coding	161	ccataacatctgcttaatat	70	18
35	32011	Coding	177	aattgtgaacttgcttccat	0	19
	32012	Coding	194	gaggttgaacattgggtaat	0	20
	32013	Coding	209	gtcaatatccataaggaggt	38	21
	32014	Coding	225	atgcaaacatataggagtca	84	22

-80-

	32015	Coding	303	ctggaagaaaaggtctgaca	55	23
	32016	Coding	408	gataccttcaaggaatcaa	49	24
	32017	Coding	544	aagttttcagggatggatg	52	25
	32018	Coding	587	atgaacagctacgatgagct	29	26
5	32019	Coding	612	taaacacgtcctggcagttt	41	27
	32020	Coding	635	catattaggagacacttgaa	77	28
	32021	Coding	662	tgccaattcatttactttga	86	29
	32022	Coding	692	cttcccatgaatagtcaa	0	30
	32023	Coding	727	tgcaacacataatcatagg	0	31
10	32024	Coding	772	attagtggatgatcaccaa	58	32
	32025	Coding	800	cacacagttccggatata	34	33
	32026	Coding	859	tacattttcttgatcttgca	54	34
	32027	Coding	1049	accagccctgacatgaactt	53	35
	32028	Coding	1360	tgtaatatattatgtctccagt	71	36
15	32029	Coding	1777	ggccaaatctgaagcagcgc	40	37
	32030	Coding	1894	tattgagaaaagttcttc	54	38
	32031	Coding	2132	attgagtgtctcaacctgct	0	39
	32032	Coding	2285	gaggataaacacatgggttca	33	40
	32033	Coding	2440	aacatttggagtgtcaacat	0	41
20	32034	Coding	2497	taaggcaacatccgaagatc	15	42
	32035	Coding	2706	agtagccagcacaggacagt	65	43
	32036	Coding	2747	actatgtctgtcaccaatcc	57	44
	32037	Coding	2975	attcccatgccgtcgtaaaa	8	45
	32038	Coding	2999	cagcgcaaagagagtgatga	26	46
25	32039	Coding	3154	ttcacttttagtagtccagct	66	47

As shown in Table 2, SEQ ID NOs 8, 9, 10, 11, 12, 13, 14, 16, 17, 18, 21, 22, 23, 24, 25, 26, 27, 28, 29, 32, 33, 34, 35, 36, 37, 38, 40, 43, 44, 46 and 47 demonstrated at

~~least 20% inhibition of PI3 kinase p110 beta expression in~~  
this experiment and are therefore preferred.

#### Example 17

#### Western blot analysis of PI3 kinase p110 beta protein levels

35 Western blot analysis (immunoblot analysis) is carried out using standard methods. Cells are harvested 16-20 h after oligonucleotide treatment, washed once with PBS, suspended in Laemmli buffer (100 ul/well), boiled for 5 minutes and loaded on a 16% SDS-PAGE gel. Gels are run for  
40 1.5 hours at 150 V, and transferred to membrane for western blotting. Appropriate primary antibody directed to PI3

-81-

kinase p110 beta is used, with a radiolabelled or fluorescently labeled secondary antibody directed against the primary antibody species. Bands are visualized using a PHOSPHORIMAGER™ (Molecular Dynamics, Sunnyvale CA).

-82-

What is claimed is:

1. An antisense compound 8 to 30 nucleobases in length targeted to a nucleic acid molecule encoding human PI3 kinase p110 beta, wherein said antisense compound specifically hybridizes with and inhibits the expression of human PI3  
5 kinase p110 beta.
2. The antisense compound of claim 1 which is an antisense oligonucleotide.
3. The antisense compound of claim 2 wherein the antisense oligonucleotide has a sequence comprising SEQ ID NO:  
10 11, 16, 19, 20, 26, 30, 35, 44, 8, 9, 10, 12, 13, 14, 17, 18, 21, 22, 23, 24, 25, 27, 28, 29, 32, 33, 34, 36, 37, 38, 40, 43, 46 or 47.
4. The antisense compound of claim 2 wherein the antisense oligonucleotide has a sequence comprising SEQ ID NO:  
15 11, 16, 26, 35 or 44.
5. The antisense compound of claim 2 wherein the antisense oligonucleotide comprises at least one modified internucleoside linkage.
6. The antisense compound of claim 5 wherein the  
20 modified internucleoside linkage is a phosphorothioate linkage.
7. The antisense compound of claim 2 wherein the antisense oligonucleotide comprises at least one modified sugar moiety.
- 25 8. The antisense compound of claim 7 wherein the modified sugar moiety is a 2'-O-methoxyethyl sugar moiety.
9. The antisense compound of claim 2 wherein the antisense oligonucleotide comprises at least one modified nucleobase.
- 30 10. The antisense compound of claim 9 wherein the modified nucleobase is a 5-methylcytosine.
11. The antisense compound of claim 2 wherein the antisense oligonucleotide is a chimeric oligonucleotide.
12. A composition comprising the antisense compound of

-83-

claim 1 and a pharmaceutically acceptable carrier or diluent.

13. The composition of claim 12 further comprising a colloidal dispersion system.

14. The composition of claim 12 wherein the antisense compound is an antisense oligonucleotide.

5        15. A method of inhibiting the expression of PI3 kinase p110 beta in human cells or tissues comprising contacting said cells or tissues with the antisense compound of claim 1 so that expression of PI3 kinase p110 beta is inhibited.

10        16. A method of treating a human having a disease or condition associated with PI3 kinase p110 beta comprising administering to said animal a therapeutically or prophylactically effective amount of the antisense compound of claim 1 so that expression of PI3 kinase p110 beta is inhibited.

15        17. The method of claim 16 wherein the disease or condition is a hyperproliferative disorder.

18. The method of claim 17 wherein the hyperproliferative disorder is cancer.

20        19. The method of claim 16 wherein the disease or condition is an inflammatory disorder.

20. The method of claim 19 wherein the inflammatory disorder is asthma or rheumatoid arthritis.

\_\_\_\_\_

## SEQUENCE LISTING

<110> Brett P. Monia  
Lex M. Cowsert  
ISIS PHARMACEUTICALS, INC.

<120> ANTISENSE MODULATION OF PI3 KINASE P110 BETA EXPRESSION

<130> RTSP-0052

<150> 09/392,530

<151> 1999-09-09

<160> 47

<210> 1

<211> 3213

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(3213)

<400> 1

atg tgc ttc agt ttc ata atg cct cct gct atg gca gac atc ctt gac	48
Met Cys Phe Ser Phe Ile Met Pro Pro Ala Met Ala Asp Ile Leu Asp	
1 5 10 15	
atc tgg gcg gtg gat tca cag ata gca tct gat ggc tcc ata cct gtg	96
Ile Trp Ala Val Asp Ser Gln Ile Ala Ser Asp Gly Ser Ile Pro Val	
20 25 30	
gat ttc ctt ttg ccc act ggg att tat atc cag ttg gag gta cct cgg	144
Asp Phe Leu Leu Pro Thr Gly Ile Tyr Ile Gln Leu Glu Val Pro Arg	
35 40 45	
gaa gct acc att tct tat att aag cag atg tta tgg aag caa gtt cac	192
Glu Ala Thr Ile Ser Tyr Ile Lys Gln Met Leu Trp Lys Gln Val His	
50 55 60	
aat tac cca atg ttc aac ctc ctt atg gat att gac tcc tat atg ttt	240
Asn Tyr Pro Met Phe Asn Leu Leu Met Asp Ile Asp Ser Tyr Met Phe	
65 70 75 80	
gca tgt gtg aat cag act gct gta tat gag gag ctt gaa gat gaa aca	288
Ala Cys Val Asn Gln Thr Ala Val Tyr Glu Glu Leu Glu Asp Glu Thr	
85 90 95	
cga aga ctc tgt gat gtc aga cct ttt ctt cca gtt ctc aaa tta gtg	336
Arg Arg Leu Cys Asp Val Arg Pro Phe Leu Pro Val Leu Lys Leu Val	
100 105 110	
aca aga agt tgt gac cca ggg gaa aaa tta gac tca aaa att gga gtc	384
Thr Arg Ser Cys Asp Pro Gly Glu Lys Leu Asp Ser Lys Ile Gly Val	
115 120 125	
ctt ata gga aaa ggt ctg cat gaa ttt gat tcc ttg aag gat cct gaa	432
Leu Ile Gly Lys Gly Leu His Glu Phe Asp Ser Leu Lys Asp Pro Glu	
130 135 140	
gta aat gaa ttt cga aga aaa atg cgc aaa ttc agc gag gaa aaa atc	480
Val Asn Glu Phe Arg Arg Lys Met Arg Lys Phe Ser Glu Glu Lys Ile	





145	150	155	160	
ctg tca ctt gtg gga ttg tct tgg atg gac tgg cta aaa caa aca tat				528
Leu Ser Leu Val Gly L u Ser Trp Met Asp Trp Leu Lys Gln Thr Tyr				
	165	170	175	
cca cca gag cat gaa cca tcc atc cct gaa aac tta gaa gat aaa ctt				576
Pro Pro Glu His Glu Pro Ser Ile Pro Glu Asn Leu Glu Asp Lys Leu				
	180	185	190	
tat ggg gga aag ctc atc gta gct gtt cat ttt gaa aac tgc cag gac				624
Tyr Gly Gly Lys Leu Ile Val Ala Val His Phe Glu Asn Cys Gln Asp				
	195	200	205	
gtg ttt agc ttt caa gtg tct cct aat atg aat cct atc aaa gta aat				672
Val Phe Ser Phe Gln Val Ser Pro Asn Met Asn Pro Ile Lys Val Asn				
	210	215	220	
gaa ttg gca atc caa aaa cgt ttg act att cat ggg aag gaa gat gaa				720
Glu Leu Ala Ile Gln Lys Arg Leu Thr Ile His Gly Lys Glu Asp Glu				
	225	230	235	240
gtt agc ccc tat gat tat gtg ttg caa gtc agc ggg aga gta gaa tat				768
Val Ser Pro Tyr Asp Tyr Val Leu Gln Val Ser Gly Arg Val Glu Tyr				
	245	250	255	
gtt ttt ggt gat cat cca cta att cag ttc cag tat atc cgg aac tgt				816
Val Phe Gly Asp His Pro Leu Ile Gln Phe Gln Tyr Ile Arg Asn Cys				
	260	265	270	
gtg atg aac aga gcc ctg ccc cat ttt ata ctt gtg gaa tgc tgc aag				864
Val Met Asn Arg Ala Leu Pro His Phe Ile Leu Val Glu Cys Cys Lys				
	275	280	285	
atc aag aaa atg tat gaa caa gaa atg att gcc ata gag gct gcc ata				912
Ile Lys Lys Met Tyr Glu Gln Glu Met Ile Ala Ile Glu Ala Ala Ile				
	290	295	300	
aat cga aat tca tct aat ctt cct ctt cca tta cca cca aag aaa aca				960
Asn Arg Asn Ser Ser Asn Leu Pro Leu Pro Leu Pro Pro Lys Lys Thr				
	305	310	315	320
cga att att tct cat gtt tgg gaa aat aac aac cct ttc caa att gtc				1008
Arg Ile Ile Ser His Val Trp Glu Asn Asn Asn Pro Phe Gln Ile Val				
	325	330	335	
ttg gtt aag gga aat aaa ctt aac aca gag gaa act gta aaa gtt cat				1056
Leu Val Lys Gly Asn Lys Leu Asn Thr Glu Glu Thr Val Lys Val His				
	340	345	350	
gtc agg gct ggt ctt ttt cat ggt act gag ctc ctg tgt aaa acc atc				1104
Val Arg Ala Gly Leu Phe His Gly Thr Glu Leu Leu Cys Lys Thr Ile				
	355	360	365	
gta agc tca gag gta tca ggg aaa aat gat cat att tgg aat gaa cca				1152
Val Ser Ser Glu Val Ser Gly Lys Asn Asp His Ile Trp Asn Glu Pro				
	370	375	380	
ctg gaa ttt gat att aat att tgt gac tta cca aga atg gct cga tta				1200
Leu Glu Phe Asp Ile Asn Ile Cys Asp Leu Pro Arg Met Ala Arg Leu				
	385	390	395	400
tgt ttt gct gtt tat gca gtt ttg gat aaa gta aaa acg aag aaa tca				1248



...



...

Cys	Phe	Ala	Val	Tyr	Ala	Val	Leu	Asp	Lys	Val	Lys	Thr	Lys	Lys	Ser		
				405					410					415			
acg	aaa	act	att	aat	ccc	tct	aaa	tat	cag	acc	atc	agg	aaa	gct	gga	1296	
Thr	Lys	Thr	Ile	Asn	Pro	Ser	Lys	Tyr	Gln	Thr	Ile	Arg	Lys	Ala	Gly		
			420					425					430				
aaa	gtg	cat	tat	cct	gta	gcg	tgg	gta	aat	acg	atg	gtt	ttt	gac	ttt	1344	
Lys	Val	His	Tyr	Pro	Val	Ala	Trp	Val	Asn	Thr	Met	Val	Phe	Asp	Phe		
		435					440					445					
aaa	gga	caa	ttg	aga	act	gga	gac	ata	ata	tta	cac	agc	tgg	tct	tca	1392	
Lys	Gly	Gln	Leu	Arg	Thr	Gly	Asp	Ile	Ile	Leu	His	Ser	Trp	Ser	Ser		
		450				455					460						
ttt	cct	gat	gaa	ctc	gaa	gaa	atg	ttg	aat	cca	atg	gga	act	gtt	caa	1440	
Phe	Pro	Asp	Glu	Leu	Glu	Glu	Met	Leu	Asn	Pro	Met	Gly	Thr	Val	Gln		
465					470					475					480		
aca	aat	cca	tat	act	gaa	aat	gca	aca	gct	ttg	cat	gtt	aaa	ttt	cca	1488	
Thr	Asn	Pro	Tyr	Thr	Glu	Asn	Ala	Thr	Ala	Leu	His	Val	Lys	Phe	Pro		
				485					490					495			
gag	aat	aaa	aaa	caa	cct	tat	tat	tac	cct	ccc	ttc	gat	aag	att	att	1536	
Glu	Asn	Lys	Lys	Gln	Pro	Tyr	Tyr	Tyr	Pro	Pro	Phe	Asp	Lys	Ile	Ile		
			500					505					510				
gaa	aag	gca	gct	gag	att	gca	agc	agt	gat	agt	gct	aat	gtg	tca	agt	1584	
Glu	Lys	Ala	Ala	Glu	Ile	Ala	Ser	Ser	Asp	Ser	Ala	Asn	Val	Ser	Ser		
		515					520					525					
cga	ggt	gga	aaa	aag	ttt	ctt	cct	gta	ttg	aaa	gaa	atc	ttg	gac	agg	1632	
Arg	Gly	Lys	Lys	Phe	Leu	Pro	Val	Leu	Lys		Glu	Ile	Leu	Asp	Arg		
	530				535						540						
gat	ccc	ttg	tct	caa	ctg	tgt	gaa	aat	gaa	atg	gat	ctt	att	tgg	act	1680	
Asp	Pro	Leu	Ser	Gln	Leu	Cys	Glu	Asn	Glu	Met	Asp	Leu	Ile	Trp	Thr		
545				550					555					560			
ttg	cga	caa	gac	tgc	cga	gag	att	ttc	cca	caa	tca	ctg	cca	aaa	tta	1728	
Leu	Arg	Gln	Asp	Cys	Arg	Glu	Ile	Phe	Pro	Gln	Ser	Leu	Pro	Lys	Leu		
			565						570					575			
ctg	ctg	tca	atc	aag	tgg	aat	aaa	ctt	gag	gat	gtt	gct	cag	ctt	cag	1776	
Leu	Leu	Ser	Ile	Lys	Trp	Asn	Lys	Leu	Glu	Asp	Val	Ala	Gln	Leu	Gln		
			580				585						590				
gcg	ctg	ctt	cag	att	tgg	cct	aaa	ctg	ccc	ccc	cgg	gag	gcc	cta	gag	1824	
Ala	Leu	Leu	Gln	Ile	Trp	Pro	Lys	Leu	Pro	Pro	Arg	Glu	Ala	Leu	Glu		
		595				600						605					
ctt	ctg	gat	ttc	aac	tat	cca	gac	cag	tac	gtt	cga	gaa	tat	gct	gta	1872	
Leu	Leu	Asp	Phe	Asn	Tyr	Pro	Asp	Gln	Tyr	Val	Arg	Glu	Tyr	Ala	Val		
		610				615					620						
ggc	tgc	ctg	cga	cag	atg	agt	gat	gaa	gaa	ctt	tct	caa	tat	ctt	tta	1920	
Gly	Cys	Leu	Arg	Gln	Met	Ser	Asp	Glu	Glu	Leu	Ser	Gln	Tyr	Leu	Leu		
		625			630					635				640			
caa	ctg	gtg	caa	gtg	tta	aaa	tat	gag	cct	ttt	ctt	gat	tgt	gcc	ctc	1968	
Gln	Leu	Val	Gln	Val	Leu	Lys	Tyr	Glu	Pro	Phe	Leu	Asp	Cys	Ala	Leu		
			645					650						655			

\_\_\_\_\_

tct aga ttc cta tta gaa aga gca ctt ggt aat cgg agg ata ggg cag Ser Arg Phe Leu Leu Glu Arg Ala Leu Gly Asn Arg Arg Ile Gly Gln 660 665 670	2016
ttt cta ttt tgg cat ctt agg tca gaa gtg cac att cct gct gtc tca Phe Leu Phe Trp His Leu Arg Ser Glu Val His Ile Pro Ala Val Ser 675 680 685	2064
gta caa ttt ggt gtc atc ctt gaa gca tac tgc cgg gga agt gtg ggg Val Gln Phe Gly Val Ile Leu Glu Ala Tyr Cys Arg Gly Ser Val Gly 690 695 700	2112
cac atg aaa gtg ctt tct aag cag gtt gaa gca ctc aat aag tta aaa His Met Lys Val Leu Ser Lys Gln Val Glu Ala Leu Asn Lys Leu Lys 705 710 715 720	2160
act tta aat agt tta atc aaa ctg aat gcc gtg aag tta aac aga gcc Thr Leu Asn Ser Leu Ile Lys Leu Asn Ala Val Lys Leu Asn Arg Ala 725 730 735	2208
aaa ggg aag gag gcc atg cat acc tgt tta aaa cag agt gct tac cgg Lys Gly Lys Glu Ala Met His Thr Cys Leu Lys Gln Ser Tyr Arg 740 745 750	2256
gaa gcc ctc tct gac ctg cag tca ccc ctg aac cca tgt gtt atc ctc Glu Ala Leu Ser Asp Leu Gln Ser Pro Leu Asn Pro Cys Val Ile Leu 755 760 765	2304
tca gaa ctc tat gtt gaa aag tgc aaa tac atg gat tcc aaa atg aag Ser Glu Leu Tyr Val Glu Lys Cys Lys Tyr Met Asp Ser Lys Met Lys 770 775 780	2352
cct ttg tgg ctg gta tac aat aac aag gta ttt ggt gag gat tca gtt Pro Leu Trp Leu Val Tyr Asn Asn Lys Val Phe Gly Glu Asp Ser Val 785 790 795 800	2400
gga gtg att ttt aaa aat ggt gat gat tta cga cag gat atg ttg aca Gly Val Ile Phe Lys Asn Gly Asp Asp Leu Arg Gln Asp Met Leu Thr 805 810 815	2448
ctc caa atg ttg cgc ttg atg gat tta ctc tgg aaa gaa gct ggt ttg Leu Gln Met Leu Arg Leu Met Asp Leu Leu Trp Lys Glu Ala Gly Leu 820 825 830	2496
gat ctt cgg atg ttg cct tat ggc tgt tta gca aca gga gat cgc tct Asp Leu Arg Met Leu Pro Tyr Gly Cys Leu Ala Thr Gly Asp Arg Ser 835 840 845	2544
ggc ctc att gaa gtt gtg agc acc tct gaa aca att gct gac att cag Gly Leu Ile Glu Val Val Ser Thr Ser Glu Thr Ile Ala Asp Ile Gln 850 855 860	2592
ctg aac agt agc aat gtg gct gct gca gca gcc ttc aac aaa gat gcc Leu Asn Ser Ser Asn Val Ala Ala Ala Ala Phe Asn Lys Asp Ala 865 870 875 880	2640
ctt ctg aac tgg ctt aaa gaa tac aac tct ggg gat gac ctg gac cga Leu Leu Asn Trp Leu Lys Glu Tyr Asn Ser Gly Asp Asp Leu Asp Arg 885 890 895	2688
gcc att gag gaa ttt aca ctg tcc tgt gct ggc tac tgt gta gct tct Ala Ile Glu Phe Thr Leu Ser Cys Ala Gly Tyr Cys Val Ala Ser 900 905 910	2736



tat gtc ctt ggg att ggt gac aga cat agt gac aac atc atg gtc aaa	2784
Tyr Val Leu Gly Ile Gly Asp Arg His Ser Asp Asn Ile Met Val Lys	
915 920 925	
aaa act ggc cag ctc ttc cac att gac ttt gga cat att ctt gga aat	2832
Lys Thr Gly Gln Leu Phe His Ile Asp Phe Gly His Ile Leu Gly Asn	
930 935 940	
ttc aaa tct aag ttt ggc att aaa agg gag cga gtg cct ttt att ctt	2880
Phe Lys Ser Lys Phe Gly Ile Lys Arg Glu Arg Val Pro Phe Ile Leu	
945 950 955 960	
acc tat gat ttc atc cat gtc att caa caa gga aaa aca gga aat aca	2928
Thr Tyr Asp Phe Ile His Val Ile Gln Gln Gly Lys Thr Gly Asn Thr	
965 970 975	
gaa aag ttt ggc cgg ttc cgc cag tgt tgt gag gat gca tat ctg att	2976
Glu Lys Phe Gly Arg Phe Arg Gln Cys Cys Glu Asp Ala Tyr Leu Ile	
980 985 990	
tta cga cgg cat ggg aat ctc ttc atc act ctc ttt gcg ctg atg ttg	3024
Leu Arg Arg His Gly Asn Leu Phe Ile Thr Leu Phe Ala Leu Met Leu	
995 1000 1005	
act gca ggg ctt cct gaa ctc aca tca gtc aaa gat ata cag tat ctt	3072
Thr Ala Gly Leu Pro Glu Leu Thr Ser Val Lys Asp Ile Gln Tyr Leu	
1010 1015 1020	
aag gac tct ctt gca tta ggg aag agt gaa gaa gaa gca ctc aaa cag	3120
Lys Asp Ser Leu Ala Leu Gly Lys Ser Glu Glu Glu Ala Leu Lys Gln	
1025 1030 1035 1040	
ttt aag caa aaa ttt gat gag gcg ctc agg gaa agc tgg act act aaa	3168
Phe Lys Gln Lys Phe Asp Glu Ala Leu Arg Glu Ser Trp Thr Thr Lys	
1045 1050 1055	
gtg aac tgg atg gcc cac aca gtt cgg aaa gac tac aga tct taa	3213
Val Asn Trp Met Ala His Thr Val Arg Lys Asp Tyr Arg Ser	
1060 1065 1070	

<210> 2  
 <211> 21  
 <212> DNA  
 <213> Artificial Sequence

<223> PCR Primer

<400> 2  
 gcgggtggatt cacagatagc a 21

<210> 3  
 <211> 21  
 <212> DNA  
 <213> Artificial Sequence

<223> PCR Primer

<400> 3  
 tggtagcttc ccgaggtacc t 21





<210> 4  
<211> 24  
<212> DNA  
<213> Artificial Sequence

<223> PCR Probe

<400> 4  
tggatttcct ttgcccact ggga

24

<210> 5  
<211> 19  
<212> DNA  
<213> Artificial Sequence

<223> PCR Primer

<400> 5  
gaaggtgaag gtcggagtc

19

<210> 6  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<223> PCR Primer

<400> 6  
gaagatggtg atgggatttc

20

<210> 7  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<223> PCR Probe

<400> 7  
caagcttccc gttctcagcc

20

<210> 8  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 8  
cattatgaaa ctgaagcaca

20

<210> 9  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 9



ggcattatga aactgaagca 20

<210> 10  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 10  
gatgtctgcc atagcaggag 20

<210> 11  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 11  
caccgcccag atgtcaagga 20

<210> 12  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 12  
tgctatctgt gaatccaccg 20

<210> 13  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 13  
aggtatggag ccatcagatg 20

<210> 14  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 14  
gggcaaaagg aaatccacag 20

<210> 15  
<211> 20  
<212> DNA  
<213> Artificial Sequence



## &lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 15

actggatata aatcccagtg

20

&lt;210&gt; 16

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

## &lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 16

ttcccagagt acctccaact

20

&lt;210&gt; 17

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

## &lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 17

tataagaaat ggtagcttcc

20

&lt;210&gt; 18

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

## &lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 18

ccataacatc tgcttaatat

20

&lt;210&gt; 19

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

## &lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 19

aattgtgaac ttgcttccat

20

&lt;210&gt; 20

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

## &lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 20

gaggttgaac attgggtaat

20

&lt;210&gt; 21

&lt;211&gt; 20



<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 21  
gtcaatatcc ataaggaggt 20

<210> 22  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 22  
atgcaaacat ataggagtca 20

<210> 23  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 23  
ctggaagaaa aggtctgaca 20

<210> 24  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 24  
gataccttcaa ggaatcaaat 20

<210> 25  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 25  
aagttttcag ggatggatgg 20

<210> 26  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 26  
atgaacagct acgatgagct 20





<210> 27  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 27  
taaacacgtc ctggcagttt 20

<210> 28  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 28  
catattagga gacacttgaa 20

<210> 29  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 29  
tgccaattca tttactttga 20

<210> 30  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 30  
cttcccatga atagtcaaac 20

<210> 31  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 31  
tgcaacacat aatcataggg 20

<210> 32  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 32

5

\_\_\_\_\_

5

5

attagtggat gatcaccaaa 20

<210> 33  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 33  
cacacagttc cggatatact 20

<210> 34  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 34  
tacattttct tgatcttgca 20

<210> 35  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 35  
accagccctg acatgaactt 20

<210> 36  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 36  
tgtaatatta tgtctccagt 20

<210> 37  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 37  
ggccaaatct gaagcagcgc 20

<210> 38  
<211> 20  
<212> DNA  
<213> Artificial Sequence



<223> Antisense Oligonucleotide

<400> 38

tattgagaaa gttcttcac

20

<210> 39

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 39

attgagtgt tcaacctgt

20

<210> 40

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 40

gaggataaca catgggttca

20

<210> 41

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 41

aacatttgga gtgtcaacat

20

<210> 42

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 42

taaggcaaca tccgaagatc

20

<210> 43

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 43

agtagccagc acaggacagt

20

<210> 44

<211> 20



6



6

<212> DNA  
<213> Artificial Sequence  
<223> Antisense Oligonucleotide

<400> 44  
actatgtctg tcaccaatcc

20

<210> 45  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
<223> Antisense Oligonucleotide

<400> 45  
attcccatgc cgtcgtaaaa

20

<210> 46  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
<223> Antisense Oligonucleotide

<400> 46  
cagcgcaaag agagtgatga

20

<210> 47  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
<223> Antisense Oligonucleotide

<400> 47  
ttcactttag tagtccagct

20





<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC(7) : C07H 21/04; C12N 15/00; C12Q 1/68; A61K 48/00 US CL : 435/8, 91.1, 91.3, 325, 375; 536, 23.1, 23.2, 24.5, 24.3, 24.31, 24.33; 514/44 According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) U.S. : 435/8, 91.1, 91.3, 325, 375; 536, 23.1, 23.2, 24.5, 24.3, 24.31, 24.33; 514/44 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) MEDLINE, BIOSIS, CAPLUS, LIFESCI, USPATFULL, EMBASE		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,776,689 A (KARIN et al.) 07 July 1998 (07.07.98) , see especially SEQ ID NO:10.	1-3
Y	MILNER et al. Selecting effective antisense reagents on combinatorial oligonucleotide arrays. Nature Biotechnology. June 1997, Vol. 15, pages 537-541, see entire document.	1-15
Y	HU et al. Cloning of a Novel, Ubiquitously Expressed Human Phosphatidylinositol 3-Kinase and Identification of Its Binding Site on p85. Mol. Cell. Biol. December 1993, Vol. 13, No. 12, pages 7677-7688, see entire document, especially Figure 1.	1, 2, 5-11
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents:	*T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *A document defining the general state of the art which is not considered to be of particular relevance *E earlier document published on or after the international filing date *L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) *O document referring to an oral disclosure, use, exhibition or other means *P document published prior to the international filing date but later than the priority date claimed *X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art *G document member of the same patent family	
Date of the actual completion of the international search 06 NOVEMBER 2000		Date of mailing of the international search report 28 DEC 2000
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3930		Authorized officer <i>Jayne Bridges for</i> KAREN A. LAPOURCIERE Telephone No. (703) 308-0196

International application No.  
PCT/US00/24388

International application No.  
PCT/US00/24388

[illegible]

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.: 16-20  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
- Claims 16-20 are drawn to in vivo, whole organism, methods of treatment and are considered to be drawn to non-statutory subject matter. Instead, claims 16-20 were searched on the alleged effects of the claimed composition.
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.



# ATENT COOPERATION TREATY

## PCT

REC'D 15 MAY 2001

WIPO


PCT

### INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)

Applicant's or agent's file reference RTSP-0052	FOR FURTHER ACTION See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/409)	
International application No. PCT/US00/24382	International filing date (day/month/year) 06 SEPTEMBER 2000	Priority date (day/month/year) 09 SEPTEMBER 1999
International Patent Classification (IPC) or national classification and IPC Please See Supplemental Sheet.		
Applicant ISIS PHARMACEUTICALS, INC.		

<p>1. This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.</p> <p>2. This REPORT consists of a total of <u>5</u> sheets.</p> <p><input type="checkbox"/> This report is also accompanied by ANNEXES, i.e., sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority. (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).</p> <p>These annexes consist of a total of <u>0</u> sheets.</p>
<p>3. This report contains indications relating to the following items:</p> <p>I <input checked="" type="checkbox"/> Basis of the report</p> <p>II <input type="checkbox"/> Priority</p> <p>III <input checked="" type="checkbox"/> Non-establishment of report with regard to novelty, inventive step or industrial applicability</p> <p>IV <input type="checkbox"/> Lack of unity of invention</p> <p>V <input checked="" type="checkbox"/> Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability, citations and explanations supporting such statement</p> <p>VI <input type="checkbox"/> Certain documents cited</p> <p>VII <input type="checkbox"/> Certain defects in the international application</p> <p>VIII <input type="checkbox"/> Certain observations on the international application</p>

Date of submission of the demand 05 APRIL 2001	Date of completion of this report 02 MAY 2001
Name and mailing address of the IPEA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231	Authorized officer  KAREN A. LACOURCIERE
Facsimile No. (703) 305-3230	Telephone No. (703) 308-0196



## INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No.

PCT/US00/24382

**I. Basis of the report**1. With regard to the **elements** of the international application:\*☒ the international application as originally filed☒ the description:pages 1-81, as originally filedpages NONE, filed with the demandpages NONE, filed with the letter of \_\_\_\_\_☒ the claims:pages 82-83, as originally filedpages NONE, as amended (together with any statement) under Article 19pages NONE, filed with the demandpages NONE, filed with the letter of \_\_\_\_\_☒ the drawings:pages NONE, as originally filedpages NONE, filed with the demandpages NONE, filed with the letter of \_\_\_\_\_☒ the sequence listing part of the description:pages 1-13, as originally filedpages NONE, filed with the demandpages NONE, filed with the letter of \_\_\_\_\_2. With regard to the **language**, all the elements marked above were available or furnished to this Authority in the language in which the international application was filed, unless otherwise indicated under this item.

These elements were available or furnished to this Authority in the following language \_\_\_\_\_ which is:

☐ the language of a translation furnished for the purposes of international search (under Rule 23.1(b)).☐ the language of publication of the international application (under Rule 48.3(b)).☐ the language of the translation furnished for the purposes of international preliminary examination (under Rules 55.2 and/or 55.3).3. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international preliminary examination was carried out on the basis of the sequence listing:☒ contained in the international application in printed form.☒ filed together with the international application in computer readable form.☒ furnished subsequently to this Authority in written form.☒ furnished subsequently to this Authority in computer readable form.☒ The statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.☒ The statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished.4. ☒ The amendments have resulted in the cancellation of:☒ the description, pages NONE☒ the claims, Nos. NONE☒ the drawings, sheets/fig NONE5. ☐ This report has been drawn as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed, as indicated in the Supplemental Box (Rule 70.2(c)).\*\*

\* Replacement sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to this report since they do not contain amendments (Rules 70.16 and 70.17).

\*\*Any replacement sheet containing such amendments must be referred to under item 1 and annexed to this report.





# INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No.  
PCT/US00/24382

## III. Non-establishment of opinion with regard to novelty, inventive step and industrial applicability

1. The questions whether the claimed invention appears to be novel, to involve an inventive step (to be non obvious), or to be industrially applicable have not been and will not be examined in respect of:

☐ the entire international application.

☒ claims Nos. 16-20 IN PART

because:

☐ the said international application, or the said claim Nos. \_ relate to the following subject matter which does not require international preliminary examination (*specify*).

☐ the description, claims or drawings (*indicate particular elements below*) or said claims Nos. \_ are so unclear that no meaningful opinion could be formed (*specify*).

☐ the claims, or said claims Nos. \_ are so inadequately supported by the description that no meaningful opinion could be formed.

☒ no international search report has been established for said claims Nos. 16-20 IN PART.

2. A meaningful international preliminary examination cannot be carried out due to the failure of the nucleotide and/or amino acid sequence listing to comply with the standard provided for in Annex C of the Administrative Instructions:

☐ the written form has not been furnished or does not comply with the standard.

☐ the computer readable form has not been furnished or does not comply with the standard.



# INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No.

PCT/US00/24382

## V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement

### 1. statement

Novelty (N)	Claims 1-20	YES
	Claims none	NO
Inventive Step (IS)	Claims 1-20	YES
	Claims none	NO
Industrial Applicability (IA)	Claims 1-20	YES
	Claims none	NO

### 2. citations and explanations (Rule 70.7)

Claims 1-20 meet the criteria set out in PCT Article 35(2)-(4), because the prior art does not teach or fairly suggest antisense of 8-30 nucleobases in length targeted to human P13kinase p110 beta with the specific sequences and modifications claimed. Nor does the prior art teach or fairly suggest the methods of treatment using said antisense, as claimed.

NEW CITATIONS

NONE



INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No.

PCT/US00/24382

**Supplemental Box**

(To be used when the space in any of the preceding boxes is not sufficient)

Continuation of Boxes I - VIII

Sheet 10


**CLASSIFICATION:**

The International Patent Classification (IPC) and/or the National classification are as listed below:

IPC(7): C07H 21/04; C12N 15/00; C12Q 1/68; A61K 48/00 and US Cl.: 435/6, 91.1, 91.3, 325, 375; 536, 23.1, 23.2, 24.5, 24.3, 24.31, 24.33; 514/44



## PATENT COOPERATION TREATY



PCT

From the INTERNATIONAL BUREAU

To:

LICATA, Jane, Massey,  
Law Offices of Jane Massey Licata  
66 E. Main Street  
Marlton, NJ 08053  
ETATS-UNIS D'AMERIQUE

NOTIFICATION CONCERNING  
SUBMISSION OR TRANSMITTAL  
OF PRIORITY DOCUMENT

(PCT Administrative Instructions, Section 411)

Date of mailing (day/month/year) 04 January 2001 (04.01.01)	<b>IMPORTANT NOTIFICATION</b>
Applicant's or agent's file reference RTSP-0052	
International application No. PCT/US00/24382	International filing date (day/month/year) 06 September 2000 (06.09.00)
International publication date (day/month/year) Not yet published	Priority date (day/month/year) 09 September 1999 (09.09.99)
Applicant ISIS PHARMACEUTICALS, INC. et al	

1. The applicant is hereby notified of the date of receipt (except where the letters "NR" appear in the right-hand column) by the International Bureau of the priority document(s) relating to the earlier application(s) indicated below. Unless otherwise indicated by an asterisk appearing next to a date of receipt, or by the letters "NR", in the right-hand column, the priority document concerned was submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b).
2. This updates and replaces any previously issued notification concerning submission or transmittal of priority documents.
3. An asterisk(\*) appearing next to a date of receipt, in the right-hand column, denotes a priority document submitted or transmitted to the International Bureau but not in compliance with Rule 17.1(a) or (b). In such a case, **the attention of the applicant is directed to Rule 17.1(c) which provides that no designated Office may disregard the priority claim concerned before giving the applicant an opportunity, upon entry into the national phase, to furnish the priority document within a time limit which is reasonable under the circumstances.**
4. The letters "NR" appearing in the right-hand column denote a priority document which was not received by the International Bureau or which the applicant did not request the receiving Office to prepare and transmit to the International Bureau, as provided by Rule 17.1(a) or (b), respectively. In such a case, **the attention of the applicant is directed to Rule 17.1(c) which provides that no designated Office may disregard the priority claim concerned before giving the applicant an opportunity, upon entry into the national phase, to furnish the priority document within a time limit which is reasonable under the circumstances.**

<u>Priority date</u>	<u>Priority application No.</u>	<u>Country or regional Office or PCT receiving Office</u>	<u>Date of receipt of priority document</u>
09 Sept 1999 (09.09.99)	09/392,350	US	04 Dec 2000 (04.12.00)

The International Bureau of WIPO  
34, chemin des Colombettes  
1211 Geneva 20, Switzerland

Facsimile No. (41-22) 740.14.35

Authorized officer

S. Mandallaz

Telephone No. (41-22) 338.83.38

\_\_\_\_\_



## PATENT COOPERATION TREATY

MAY 16 2001

From the  
INTERNATIONAL PRELIMINARY EXAMINING AUTHORITY

PCT

NOTIFICATION OF TRANSMITTAL OF  
INTERNATIONAL PRELIMINARY  
EXAMINATION REPORT

(PCT Rule 71.1)

To: JANE MASSEY LICATA  
LICATA & TYRRELL P.C.  
66 E. MAIN STREET  
MARLTON, NJ 08053Docket System ☒  
Status Report ☒  
Docket Book ☒

np = 3/9/02

Date of Mailing  
(day/month/year)

11 MAY 2001

Applicant's or agent's file reference

RTSP-0052

## IMPORTANT NOTIFICATION

International application No.

PCT/US00/24582

International filing date (day/month/year)

06 SEPTEMBER 2000

Priority Date (day/month/year)

09 SEPTEMBER 1999

Applicant

ISIS PHARMACEUTICALS, INC.

1. The applicant is hereby notified that this International Preliminary Examining Authority transmits herewith the international preliminary examination report and its annexes, if any, established on the international
2. A copy of the report and its annexes, if any, is being transmitted to the International Bureau for communication to all the elected Offices.
3. Where required by any of the elected Offices, the International Bureau will prepare an English translation of the report (but not of any annexes) and will transmit such translation to those Offices.
4. REMINDER

The applicant must enter the national phase before each elected Office by performing certain acts (filing translations and paying national fees) within 30 months from the priority date (or later in some Offices) (Article 39(1)) (see also the reminder sent by the International Bureau with Form PCT/IB/301).

Where a translation of the international application must be furnished to an elected Office, that translation must contain a translation of any annexes to the international preliminary examination report. It is the applicant's responsibility to prepare and furnish such translation directly to each elected Office concerned.

For further details on the applicable time limits and requirements of the elected Offices, see Volume II of the PCT Applicant's Guide.

Name and mailing address of the IPEA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

KAREN A. LACOURCIERE

Telephone No. (703) 308-0196

\_\_\_\_\_

# PATENT COOPERATION TREATY

From the  
INTERNATIONAL PRELIMINARY EXAMINING AUTHORITY

To: JANE MASSEY LICATA  
LICATA & TYRRELL P.C.  
66 E. MAIN STREET  
MARLTON, NJ 08053

## PCT

### NOTIFICATION OF TRANSMITTAL OF INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Rule 71.1)

Date of Mailing  
(day/month/year)

11 MAY 2001

Applicant's or agent's file reference  
RTSP-0058

#### IMPORTANT NOTIFICATION

International application No.

PCT/US00/24388

International filing date (day/month/year)

06 SEPTEMBER 2000

Priority Date (day/month/year)

09 SEPTEMBER 1999

Applicant

ISIS PHARMACEUTICALS, INC.

1. The applicant is hereby notified that this International Preliminary Examining Authority transmits herewith the international preliminary examination report and its annexes, if any, established on the international
2. A copy of the report and its annexes, if any, is being transmitted to the International Bureau for communication to all the elected Offices.
3. Where required by any of the elected Offices, the International Bureau will prepare an English translation of the report (but not of any annexes) and will transmit such translation to those Offices.
4. REMINDER

The applicant must enter the national phase before each elected Office by performing certain acts (filing translations and paying national fees) within 30 months from the priority date (or later in some Offices) (Article 39(1)) (see also the reminder sent by the International Bureau with Form PCT/IB/301).

Where a translation of the international application must be furnished to an elected Office, that translation must contain a translation of any annexes to the international preliminary examination report. It is the applicant's responsibility to prepare and furnish such translation directly to each elected Office concerned.

For further details on the applicable time limits and requirements of the elected Offices, see Volume II of the PCT Applicant's Guide.

Name and mailing address of the IPEA/US

Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

KAREN A. LACOURTIERE

Telephone No. (703) 308-0196



## PATENT COOPERATION TREATY

## PCT

## INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)

Applicant's or agent's file reference RTSP-0052	<b>FOR FURTHER ACTION</b> See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/409)	
International application No. PCT/US00/24382	International filing date (day/month/year) 06 SEPTEMBER 2000	Priority date (day/month/year) 09 SEPTEMBER 1999
International Patent Classification (IPC) or national classification and IPC Please See Supplemental Sheet.		
Applicant ISIS PHARMACEUTICALS, INC.		

1. This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.

2. This REPORT consists of a total of 5 sheets.

☐ This report is also accompanied by ANNEXES, i.e., sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority. (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).

These annexes consist of a total of 0 sheets.

3. This report contains indications relating to the following items:

I ☒ Basis of the report

II ☐ Priority

III ☒ Non-establishment of report with regard to novelty, inventive step or industrial applicability


IV ☐ Lack of unity of invention

V ☒ Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement

VI ☐ Certain documents cited

VII ☐ Certain defects in the international application

VIII ☐ Certain observations on the international application

Date of submission of the demand  05 APRIL 2001	Date of completion of this report  02 MAY 2001
Name and mailing address of the IPEA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231	Authorized officer  KAREN A. LACOURCIERE
Facsimile No. (703) 305-3830	Telephone No. (703) 308-0196



## INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No.

PCT/US00/24382

**I. Basis of the report****1. With regard to the elements of the international application:\***☒ the international application as originally filed☒ the description:

pages 1-81

, as originally filed

pages NONE

, filed with the demand

pages NONE

, filed with the letter of

☒ the claims:

pages 82-83

, as originally filed

pages NONE

, as amended (together with any statement) under Article 19

pages NONE

, filed with the demand

pages NONE

, filed with the letter of

☒ the drawings:

pages NONE

, as originally filed

pages NONE

, filed with the demand

pages NONE

, filed with the letter of

☒ the sequence listing part of the description:

pages 1-13

, as originally filed

pages NONE

, filed with the demand

pages NONE

, filed with the letter of

**2. With regard to the language, all the elements marked above were available or furnished to this Authority in the language in which the international application was filed, unless otherwise indicated under this item.**

These elements were available or furnished to this Authority in the following language \_\_\_\_\_ which is:

☐ the language of a translation furnished for the purposes of international search (under Rule 23.1(b)).☐ the language of publication of the international application (under Rule 48.3(b)).☐ the language of the translation furnished for the purposes of international preliminary examination (under Rules 55.2 and/or 55.3).**3. With regard to any nucleotide and/or amino acid sequence disclosed in the international application, the international preliminary examination was carried out on the basis of the sequence listing:**☒ contained in the international application in printed form.☒ filed together with the international application in computer readable form.☒ furnished subsequently to this Authority in written form.☒ furnished subsequently to this Authority in computer readable form.☒ The statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.☒ The statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished.**4. ☒ The amendments have resulted in the cancellation of:**☒ the description, pages NONE☒ the claims, Nos. NONE☒ the drawings, sheets/fig. NONE**5. ☐ This report has been drawn as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed, as indicated in the Supplemental Box (Rule 70.2(c)).\*\***

\* Replacement sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to this report since they do not contain amendments (Rules 70.16 and 70.17).

\*\*Any replacement sheet containing such amendments must be referred to under item 1 and annexed to this report.





INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No.  
PCT/US00/24382

III. Non-establishment of opinion with regard to novelty, inventive step and industrial applicability

1. The questions whether the claimed invention appears to be novel, to involve an inventive step (to be non obvious), or to be industrially applicable have not been and will not be examined in respect of:

☐ the entire international application.

☒ claims Nos. 16-20 IN PART

because:

☐ the said international application, or the said claim Nos. \_ relate to the following subject matter which does not require international preliminary examination (*specify*).

☐ the description, claims or drawings (*indicate particular elements below*) or said claims Nos. \_ are so unclear that no meaningful opinion could be formed (*specify*).

☐ the claims, or said claims Nos. \_ are so inadequately supported by the description that no meaningful opinion could be formed.

☒ no international search report has been established for said claims Nos. 16-20 IN PART.

2. A meaningful international preliminary examination cannot be carried out due to the failure of the nucleotide and/or amino acid sequence listing to comply with the standard provided for in Annex C of the Administrative Instructions:

☐ the written form has not been furnished or does not comply with the standard.

☐ the computer readable form has not been furnished or does not comply with the standard.

\_\_\_\_\_

# INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No.

PCT/US00/24382

## V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement

### 1. statement

Novelty (N)	Claims 1-20	YES
	Claims none	NO
Inventive Step (IS)	Claims 1-20	YES
	Claims none	NO
Industrial Applicability (IA)	Claims 1-20	YES
	Claims none	NO

### 2. citations and explanations (Rule 70.7)

Claims 1-20 meet the criteria set out in PCT Article 33(2)-(4), because the prior art does not teach or fairly suggest antisense of 8-30 nucleobases in length targeted to human P13kinase p110 beta with the specific sequences and modifications claimed. Nor does the prior art teach or fairly suggest the methods of treatment using said antisense, as claimed.

NEW CITATIONS

NONE

---

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No.

PCT/US00/24382

**Supplemental Box**

(To be used when the space in any of the preceding boxes is not sufficient)

Continuation of: Boxes I - VIII

Sheet 10

**CLASSIFICATION:**

The International Patent Classification (IPC) and/or the National classification are as listed below:

IPC(7): C07H 21/04; C12N 15/00; C12Q 1/68; A61K 48/00 and US Cl.: 435/6, 91.1, 91.3, 325, 375; 536, 23.1, 23.2, 24.5, 24.3, 24.31, 24.33; 514/44

\_\_\_\_\_

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/24382

**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(7) : C07H 21/04; C12N 15/00; C12Q 1/68; A61K 48/00

US CL : 435/6, 91.1, 91.3, 325, 375; 536, 23.1, 23.2, 24.5, 24.3, 24.31, 24.33; 514/44

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/6, 91.1, 91.3, 325, 375; 536, 23.1, 23.2, 24.5, 24.3, 24.31, 24.33; 514/44

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

MEDLINE, BIOSIS, CAPLUS, LIFESCI, USPATFULL, EMBASE

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,776,689 A (KARIN et al.) 07 July 1998 (07.07.98) , see especially SEQ ID NO:10.	1-3
Y	MILNER et al. Selecting effective antisense reagents on combinatorial oligonucleotide arrays. Nature Biotechnology. June 1997, Vol. 15, pages 537-541, see entire document.	1-15
Y	HU et al. Cloning of a Novel, Ubiquitously Expressed Human Phosphatidylinositol 3-Kinase and Identification of Its Binding Site on p85. Mol. Cell. Biol. December 1993, Vol. 13, No. 12, pages 7677-7688, see entire document, especially Figure 1.	1, 2, 5-11

☒ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* Special categories of cited documents:	*T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A document defining the general state of the art which is not considered to be of particular relevance	*X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*E earlier document published on or after the international filing date	*Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G document member of the same patent family
*O document referring to an oral disclosure, use, exhibition or other means	
*P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search 06 NOVEMBER 2000	Date of mailing of the international search report 28 DEC 2000
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-5230	Authorized officer <i>Karen A. LaCourciere</i> KAREN A. LA-COURCIERE Telephone No. (703) 308-0196





## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/24382

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	VLAHOS et al. A Specific Inhibitor of Phosphatidylinositol 3-Kinase, 2-(4-Morpholinyl)-8-phenyl-4H-1-benzopyran-4-one (LY294002). Journal of Biological Chemistry. 18 February 1994, Vol. 269, No. 7, Pages 5241-5248, see entire document.	1, 2, 5-11
Y	US 5,801,154 A (BARACCHINI et al.) 01 September 1998 (01.09.98), see entire document.	1,2, 5-11



..



# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/24382

## Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
  
2. ☒ Claims Nos.: 16-20  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:  
  
Claims 16-20 are drawn to in vivo, whole organism, methods of treatment and are considered to be drawn to non-statutory subject matter. Instead, claims 16-20 were searched on the alleged effects of the claimed composition.
  
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
  
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
  
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
  
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.



11

12

13

14



15  
16  
17